

**WEST****Freeform Search****Database:**

US Patents Full-Text Database  
US Pre-Grant Publication Full-Text Database  
JPO Abstracts Database  
EPO Abstracts Database  
Derwent World Patents Index  
IBM Technical Disclosure Bulletins

**Term:**

114 and (12 or 13) not 14

**Display:**

20

**Documents in Display Format:**

CIT

**Starting with Number**

1

**Generate:** ☐ Hit List ☒ Hit Count ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

**Search History****Today's Date:** 1/26/2002

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT	l14 and (l2 or l3) not l4	11	<u>L15</u>
USPT	((435/155)!.CCLS. )	188	<u>L14</u>
JPAB,EPAB,DWPI	l12 not l10	26	<u>L13</u>
JPAB,EPAB,DWPI	l8 and recombinant	34	<u>L12</u>
JPAB,EPAB,DWPI	propanediol and plasmid	1	<u>L11</u>
JPAB,EPAB,DWPI	(diol or glycerol) near2 (dehydrase or dehydratase)	21	<u>L10</u>
JPAB,EPAB,DWPI	dha\$5 and propanediol	1	<u>L9</u>
JPAB,EPAB,DWPI	dha\$5 or propanediol	5534	<u>L8</u>
USPT	(diol or glycerol) near2 (dehydrase or dehydratase)	27	<u>L7</u>
USPT	(l3 same l2) not l5	3	<u>L6</u>
USPT	l3 with l2	9	<u>L5</u>
USPT	l1 and (l2 or l3)	26	<u>L4</u>
USPT	propanediol	17930	<u>L3</u>
USPT	dha\$5	4012	<u>L2</u>
USPT	((435/158)!.CCLS. )	111	<u>L1</u>

**WEST****Freeform Search****Database:**

US Patents Full-Text Database  
 US Pre-Grant Publication Full-Text Database  
**JPO Abstracts Database**  
 EPO Abstracts Database  
 Derwent World Patents Index  
 IBM Technical Disclosure Bulletins

**Term:**

112 not 110

**Display:**

20

**Documents in Display Format:**

CIT

**Starting with Number**

1

**Generate:** ☐ Hit List ☒ Hit Count ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

**Search History****Today's Date:** 1/26/2002

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
JPAB,EPAB,DWPI	112 not 110	26	<u>L13</u>
JPAB,EPAB,DWPI	18 and recombinant	34	<u>L12</u>
JPAB,EPAB,DWPI	propanediol and plasmid	1	<u>L11</u>
JPAB,EPAB,DWPI	(diol or glycerol) near2 (dehydrase or dehydratase)	21	<u>L10</u>
JPAB,EPAB,DWPI	dha\$5 and propanediol	1	<u>L9</u>
JPAB,EPAB,DWPI	dha\$5 or propanediol	5534	<u>L8</u>
USPT	(diol or glycerol) near2 (dehydrase or dehydratase)	27	<u>L7</u>
USPT	(13 same 12) not 15	3	<u>L6</u>
USPT	13 with 12	9	<u>L5</u>
USPT	11 and (12 or 13)	26	<u>L4</u>
USPT	propanediol	17930	<u>L3</u>
USPT	dha\$5	4012	<u>L2</u>
USPT	((435/158 )!.CCLS. )	111	<u>L1</u>

**OCR WEST**[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
(diol or glycerol) near2 (dehydrase or dehydratase)	0

**Database:** [US OCR Full-Text Database](#)**Refine Search:**

(diol or glycerol) near2 (dehydrase or dehydratase)

[Clear](#)**Search History****Today's Date: 1/26/2002**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USOC	(diol or glycerol) near2 (dehydrase or dehydratase)	0	<a href="#">L7</a>
USOC	(l3 same l2 ) not l5	2	<a href="#">L6</a>
USOC	l3 with l2	2	<a href="#">L5</a>
USOC	l1 and (l2 or l3)	0	<a href="#">L4</a>
USOC	propanediol	3470	<a href="#">L3</a>
USOC	dha\$5	20237	<a href="#">L2</a>
USOC	((435/158 )!.CCLS. )	21	<a href="#">L1</a>



Generate Collection

L7: Entry 2 of 27

File: USPT

Oct 2, 2001

DOCUMENT-IDENTIFIER: US 6297428 B1

TITLE: Method for inducing viral resistance into a plant

## DEPR:

The plasmid pET-P15 (harbouring the P15 nucleic acid sequence) was restricted at its single BamHI site and blunt-ended with T4 DNA polymerase. After purification by electrophoresis in 0.8% agarose gel, the linear plasmid was restricted at its single NcoI site. The P15 gene fragment of 400 bp was purified by electrophoresis and inserted into pMJBX-Ub (harbouring the Arabidopsis polyubiquitin promoter (Norris et al., Plant Molecular Biology 21, pp. 895-906 (1993), a TMV enhancer sequence and the Nos 3' terminator) cut with NcoI and SmaI restriction endonucleases. In the plasmid so obtained (pMJBX-Ub-P15), the nucleic acid sequence of the P15 gene is placed under the control of the Arabidopsis polyubiquitin promoter followed by the TMV enhancer sequence. The EcoRI fragment from plasmid pB235SAck contains the pat gene, used as the selective marker, encoding phosphinothricin acetyl transferase (obtained from Agrevo, Berlin Germany). On this EcoRI fragment, the nucleic acid sequence of the pat gene is under the control of the 5' and 3' expression signals of the Cauliflower virus. The plasmid pMJBS6, resulting from the combination of this EcoRI-pat fragment and a partial EcoRI digestion of plasmid pMJBX-Ub-P15, contains both the pat and the P15 genes. This pMJBS6 plasmid is a high-copy plasmid based on the pUC18 vector and contains also the -lactamase gene (amp.sup.r). In the plasmid pIGPD7, harbouring the same pat fragment as pB235SAck, the -lactamase gene was replaced by an igpd (imidazole glycerol phosphate dehydratase) gene from *Saccharomyces cerevisiae* (Struhl et al., Proceedings of the National Academy of Science USA 73, pp. 1471-1475 (1976). Selection for and maintenance of the plasmid in *Escherichia coli* was achieved by complementation of an auxotrophic *hisB* strain SB3930 on minimal medium in the absence of antibiotics. The P15 fragment, with its ubiquitin promoter and terminator sequence, was purified as a 2500 bp fragment obtained from the pMJBX-Ub-P15 plasmid after it was cut at the single HindIII site, followed by a partial EcoRI restriction. This fragment was blunt-ended and inserted in a blunt-ended pIGPD7 plasmid, cut at the single NcoI site. The resulting pIGPDS4 plasmid contains both the pat and the P15 genes on a vector without the .beta.-lactamase gene.

**WEST**☐ Generate Collection

L2: Entry 5 of 6

File: USPT

Jan 11, 2000

US-PAT-NO: 6013494

DOCUMENT-IDENTIFIER: US 6013494 A

TITLE: Method for the production of 1,3-propanediol by recombinant microorganisms

DATE-ISSUED: January 11, 2000

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Nakamura; Charles E.	Claymont	DE		
Gatenby; Anthony A.	Wilmington	DE		
Hsu; Amy Kuang-Hua	Redwood City	CA		
La Reau; Richard D.	Mountain View	CA		
Haynie; Sharon L.	Philadelphia	PA		
Diaz-Torres; Maria	San Mateo	CA		
Trimbur; Donald E.	Redwood City	CA		
Whited; Gregory M.	Belmont	CA		
Nagarajan; Vasantha	Wilmington	DE		
Payne; Mark S.	Wilmington	DE		
Picataggio; Stephen K.	Landenberg	PA		
Nair; Ramesh V.	Wilmington	DE		

US-CL-CURRENT: 435/158; 435/252.3, 435/252.33, 435/254.21,  
435/69.1

## CLAIMS:

What is claimed is:

1. A method for the production of 1,3-propanediol from a recombinant microorganism comprising:
  - (i) transforming a suitable host microorganism with one or more transformation cassettes each of which comprises at least one of
    - (a) a gene encoding a glycerol-3-phosphate dehydrogenase activity;
    - (b) a gene encoding a glycerol-3-phosphatase activity;
    - (c) genes encoding a dehydratase activity; and
    - (d) a gene encoding 1,3-propanediol oxidoreductase activity,wherein all of the genes of (a)-(d) are introduced into the host microorganism;
  - (ii) culturing the transformed host microorganism under suitable conditions in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, or a one-carbon substrate whereby 1,3-propanediol

- is produced; and
- (iii) recovering the 1,3-propanediol.
2. The method of claim 1 wherein the suitable host microorganism is selected from the group consisting of bacteria, yeast, and filamentous fungi.
3. The method of claim 2 wherein the suitable host microorganism is selected from the group of genera consisting of *Citrobacter*, *Enterobacter*, *Clostridium*, *Klebsiella*, *Aerobacter*, *Lactobacillus*, *Aspergillus*, *Saccharomyces*, *Schizosaccharomyces*, *Zygosaccharomyces*, *Pichia*, *Kluyveromyces*, *Candida*, *Hansenula*, *Debaryomyces*, *Mucor*, *Torulopsis*, *Methylobacter*, *Escherichia*, *Salmonella*, *Bacillus*, *Streptomyces* and *Pseudomonas*.
4. The method of claim 3 wherein the suitable host microorganism is selected from the group consisting of *E. coli*, *Klebsiella* spp., and *Saccharomyces* spp.
5. The method of claim 1 wherein the transformed host microorganism is a *Klebsiella* spp. transformed with a transformation cassette comprising the genes GPD1 and GPP2.
6. The method claim 1 wherein the carbon source is glucose.
7. The method of claim 1 wherein the gene encoding a glycerol-3-phosphate dehydrogenase activity is selected from the group consisting of
- (a) an isolated nucleic acid molecule encoding the amino acid sequence set forth in SEQ ID NO:11, in SEQ ID NO:12, and in SEQ ID NO:13, or an enzymatically active fragment thereof;
  - (b) an isolated nucleic acid molecule that hybridizes with (a) under the following hybridization conditions: 0.1.times.SSC, 0.1% SDS at 65.degree. C.; and
  - (c) an isolated nucleic acid molecule that is completely complementary to (a) or (b).
8. The method of claim 1 wherein the gene encoding a glycerol-3-phosphatase activity is selected from the group consisting of
- (a) an isolated nucleic acid molecule encoding the amino acid sequence set forth in SEQ ID NO:33 and in SEQ ID NO:17, or an enzymatically active fragment thereof;
  - (b) an isolated nucleic acid molecule that hybridizes with (a) under the following hybridization conditions: 0.1.times.SSC, 0.1% SDS at 65.degree. C.; and
  - (c) an isolated nucleic acid molecule that is completely complementary to (a) or (b).
9. The method of claim 1 wherein the gene encoding a glycerol-3-phosphatase activity is a glycerol kinase gene selected from the group consisting of
- (a) an isolated nucleic acid molecule encoding the amino acid sequence set forth in SEQ ID NO:18, or an enzymatically active fragment thereof;
  - (b) an isolated nucleic acid molecule that hybridizes with (a) under the following hybridization conditions: 0.1.times.SSC, 0.1% SDS at 65.degree. C.; and
  - (c) an isolated nucleic acid molecule that is completely complementary to (a) or (b).
10. The method of claim 1 wherein the genes encoding a dehydratase activity comprise *dhaB1*, *dhaB2* and *dhaB3*, and are selected from the group consisting of
- (a) an isolated nucleic acid molecule encoding the amino acid sequence set forth in SEQ ID NO:34, SEQ ID NO:35, and SEQ ID NO:36, or an enzymatically active fragment thereof;

(b) an isolated nucleic acid molecule that hybridizes with (a) under the following hybridization conditions: 0.1.times.SSC, 0.1% SDS at 65.degree. C.; and (c) an isolated nucleic acid molecule that is completely complementary to (a) or (b).

11. The method of claim 1 wherein the gene encoding a 1,3-propanediol oxidoreductase activity selected from the group consisting of

(a) an isolated nucleic acid molecule encoding the amino acid sequence set forth in SEQ ID NO:37, or an enzymatically active fragment thereof;

(b) an isolated nucleic acid molecule that hybridizes with (a) under the following hybridization conditions: 0.1.times.SSC, 0.1% SDS at 65.degree. C.; and

(c) an isolated nucleic acid molecule that is completely complementary to (a) or (b).

12. A method for the production of 1,3-propanediol from a recombinant microorganism comprising:

(i) culturing, under suitable conditions in the presence of at least one carbon source selected from the group consisting of monosaccharides, oligosaccharides, polysaccharides, or a one-carbon substrate, a transformed host microorganism comprising (a) a gene encoding a glycerol-3-phosphate dehydrogenase activity;

(b) a gene encoding a glycerol-3-phosphatase activity;

(c) genes encoding a dehydratase activity; and

(d) a gene encoding 1,3-propanediol oxidoreductase activity, wherein all of the genes (a)-(d) are exogenous to the host microorganism, whereby 1,3-propanediol is produced; and

(ii) recovering the 1,3-propanediol.

13. A host cell transformed with a group of genes comprising:

(1) a gene encoding a glycerol-3-phosphate dehydrogenase enzyme corresponding to the amino acid sequence given in SEQ ID NO:11;

(2) a gene encoding a glycerol-3-phosphatase enzyme corresponding to the amino acid sequence given in SEQ ID NO:17;

(3) a gene encoding the  $\alpha$  subunit of the glycerol dehydratase enzyme corresponding to the amino acid sequence given in SEQ ID NO:34;

(4) a gene encoding the  $\beta$  subunit of the glycerol dehydratase enzyme corresponding to the amino acid sequence given in SEQ ID NO:35;

(5) a gene encoding the  $\gamma$  subunit of the glycerol dehydratase enzyme corresponding to the amino acid sequence given in SEQ ID NO:36; and

(6) a gene encoding the 1,3-propanediol oxidoreductase enzyme corresponding to the amino acid sequence given in SEQ ID NO:37, whereby the transformed host cell produces 1,3-propanediol on at least one substrate selected from the group consisting of monosaccharides, oligosaccharides, and polysaccharides or from a one-carbon substrate.



**WEST****End of Result Set**

Generate Collection

L2: Entry 6 of 6

File: USPT

Nov 11, 1997

US-PAT-NO: 5686276DOCUMENT-IDENTIFIER: US 5686276 A

TITLE: Bioconversion of a fermentable carbon source to  
1,3-propanediol by a single microorganism

DATE-ISSUED: November 11, 1997

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Laffend; Lisa Anne	Wilmington	DE		
Nagarajan; Vasantha	Wilmington	DE		
Nakamura; Charles Edwin	Claymont	DE		

US-CL-CURRENT: 435/158; 435/252.31, 435/252.33

## CLAIMS:

What is claimed is:

1. A process comprising the bioconversion of a carbon substrate, other than glycerol or dihydroxyacetone, to 1,3-propanediol by a single microorganism having at least one gene that expresses a dehydratase enzyme by contacting said microorganism with said substrate.
2. The process of claim 1 wherein said microorganism has been genetically altered.
3. The process of claim 1 wherein the dehydratase enzyme is a glycerol dehydratase enzyme or a diol dehydratase enzyme.
4. The process of claim 1 wherein the microorganism is selected from the group consisting of members of the genera Citrobacter, Enterobacter, Clostridium, Klebsiella, Aerobacter, Lactobacillus, Aspergillus, Saccharomyces, Zygosaccharomyces, Pichia, Kluyveromyces, Candida, Hansenula, Debaryomyces, Mucor, Torulopsis, Methylobacteria, Escherichia, and Salmonella; recombinant microorganisms transformed with a gene encoding a glycerol dehydratase enzyme or a diol dehydratase enzyme; and mutants of microorganisms having phenotypes which enhance production of 1,3-propanediol.
5. The process of claim 4 wherein the microorganism is selected from the group consisting of members of the genera Klebsiella and Citrobacter, and recombinant Escherichia.
6. The process of claim 5 wherein the microorganism is recombinant E. coli.
7. The process of claim 1 wherein the carbon substrate is selected from the group consisting of compounds having at least a single

carbon atom, provided that the substrate is other than glycerol or dihydroxyacetone.

8. The process of claim 7 wherein the carbon substrate is selected from the group consisting of monosaccharides and oligosaccharides.

9. The process of claim 8 wherein the carbon substrate is glucose.

10. The process of claim 1 wherein the gene is a glycerol dehydratase gene isolated from the group consisting of members of the genera *Klebsiella*, *Citrobacter*, and *Clostridium*.

11. The process of claim 1 wherein the gene is a diol dehydratase gene isolated from the group consisting of members of the genera *Klebsiella* and *Salmonella*.

12. The process of claim 1 or 9 wherein the microorganism is *E. coli* containing a glycerol dehydratase gene from *Klebsiella pneumoniae*.

13. The process of claim 1 wherein the microorganism is grown in a medium prior to contacting it with the carbon substrate.

14. A process for the bioconversion of a carbon substrate to 1,3-propanediol by a single microorganism comprising:

(i) contacting a medium containing at least one carbon substrate with a single microorganism to yield a culture medium, wherein the at least one carbon substrate is selected from the group consisting of monosaccharides, oligosaccharides, and polysaccharides, provided that the carbon substrate is other than glycerol or dihydroxyacetone, and wherein said single microorganism is selected from the group consisting of members of the genera *Klebsiella*, *Citrobacter*, recombinant *Escherichia*, or is a recombinant organism transformed with a gene encoding a diol dehydratase enzyme or a glycerol dehydratase enzyme,  
(ii) incubating said culture medium under suitable conditions to produce 1,3-propanediol; and

(iii) recovering said 1,3-propanediol.

15. The process of claim 14 wherein the at least one carbon substrate is glucose and wherein said single microorganism is a recombinant *E. coli* transformed with a gene encoding a diol dehydratase enzyme or a glycerol dehydratase enzyme.

16. The process of claim 1 further comprising recovering 1,3-propanediol following the bioconversion of the carbon substrate.

**WEST**

Generate Collection

L2: Entry 4 of 6

File: USPT

Feb 15, 2000

US-PAT-NO: 6025184

DOCUMENT-IDENTIFIER: US 6025184 A

TITLE: Bioconversion of a fermentable carbon source to  
1,3-propanediol by a single microorganism

DATE-ISSUED: February 15, 2000

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Laffend; Lisa Anne	Wilmington	DE		
Nagarajan; Vasantha	Wilmington	DE		
Nakamura; Charles Edwin	Claymont	DE		

US-CL-CURRENT: 435/252.33; 435/252.3, 435/320.1

## CLAIMS:

What is claimed is:

1. A cosmid contained in ATCC 69789 comprising a DNA fragment of about 35 kb isolated from *Klebsiella pneumoniae* wherein 1) the DNA fragment encodes an active glycerol dehydratase enzyme and 2) digestion of the cosmid results in a restriction digest pattern as shown in FIG. 1, columns 1 and 2.
2. A host bacterium transformed with the cosmid of claim 1.
3. The host bacterium of claim 2 which is deposited with the American Type Culture Collection and having accession number ATCC 69789.
4. A host bacterium comprising the cosmid of claim 1, wherein at least one DNA fragment of said cosmid encodes 1,3-propanediol oxidoreductase, and wherein said host converts a carbon source, other than glycerol or dihydroxyacetone, to 1,3-propanediol.

**WEST**☐ Generate Collection

L1: Entry 2 of 3

File: USPT

Oct 13, 1998

US-PAT-NO: 5821092

DOCUMENT-IDENTIFIER: US 5821092 A

TITLE: Production of 1,3-propanediol from glycerol by recombinant bacteria expressing recombinant diol dehydratase

DATE-ISSUED: October 13, 1998

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Nagarajan; Vasantha	Wilmington	DE		
Nakamura; Charles Edwin	Claymont	DE		

US-CL-CURRENT: 435/158; 435/232, 435/252.3, 435/252.31,  
435/252.33, 435/252.35, 435/252.5, 435/252.7, 435/320.1, 536/23.1,  
536/23.2, 536/23.7

## CLAIMS:

What is claimed is:

1. A process for the bioconversion of a carbon substrate for diol dehydratase enzyme to the corresponding product comprising the steps of:

(i) transforming a microbial host with genes encoding an enzymatically active bacterial diol dehydratase enzyme, the genes derived from

(1) a cosmid, the cosmid comprising a DNA fragment of about 35 kb isolated from *Klebsiella pneumoniae* and contained within transformed *E. coli* deposited with the American Type Culture Collection under accession number ATCC 69790; or from

(2) enzymatically active diol dehydratase genes isolated from the group consisting of members of the species *Klebsiella* sp., *Clostridia* sp., *Salmonella* sp. and *Citrobacter* sp, one subunit of the genes and having at least a 95% identity to the nucleic acid sequence of SEQ ID NO:1;

(ii) contacting the transformed microbial host with the carbon substrate in a suitable medium; and

(iii) recovering the corresponding product from the suitable medium.

2. The process of claim 1 wherein the carbon substrate is selected from the group consisting of ethylene glycol, 1,2-propanediol, glycerol and 2,3-butanediol.

3. The process of claim 2 wherein the carbon substrate is glycerol.

4. The process of claim 3 wherein the glycerol is converted to 1,3-propanediol.

**WEST****End of Result Set**☐ **Generate Collection**

L1: Entry 3 of 3

File: USPT

May 27, 1997

US-PAT-NO: 5633362DOCUMENT-IDENTIFIER: US 5633362 A

TITLE: Production of 1,3-propanediol from glycerol by recombinant bacteria expressing recombinant diol dehydratase

DATE-ISSUED: May 27, 1997

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Nagarajan; Vasantha	Wilmington	DE		
Nakamura; Charles E.	Claymont	DE		

US-CL-CURRENT: 536/23.1; 435/252.3, 435/252.33, 536/22.1, 536/24.3

## CLAIMS:

What is claimed is:

1. A cosmid comprising a DNA fragment of about 35 kb isolated from *Klebsiella pneumoniae* wherein said fragment encodes an active diol dehydratase enzyme having the restriction digest in FIG. 5, columns numbered 4, said cosmid contained within a transformed *E. coli* deposited with the American Type Culture Collection under accession number ATCC 69790.
2. A transformed microorganism comprising a host microorganism and the cosmid of claim 1.
3. The transformed microorganism of claim 2 wherein the host microorganism is *E. coli*, and which is deposited with the American Type Culture Collection as accession number ATCC 69790.
4. The cosmid of claim 1 which when transformed into bacteria causes metabolism of glycerol to 1,3-propanediol.
5. A transformed microorganism comprising a host microorganism and a DNA fragment of the cosmid of claim 1, said fragment encoding an active functional protein.
6. A DNA fragment comprising a gene encoding a diol dehydratase enzyme, said gene encompassed by the cosmid of claim 1.
7. A isolated gene encoding an active diol dehydratase enzyme comprising a contiguous sequence which consists of SEQ ID NO: 1.
8. A isolated gene encoding an active alcohol dehydrogenase comprising a contiguous sequence which consists of SEQ ID NO: 2.
9. A transformed microorganism comprising a host microorganism and the heterologous gene of claim 7 or claim 8.
10. A transformed microorganism comprising *E. coli* DH5.alpha. and the DNA sequence of claim 7 or claim 8.

5. The process of claim 1 wherein the microbial host is selected from the group consisting of members of the genera *Eschericia*, *Bacillus*, *Klebsiella*, *Citrobacter*, *Saccharomyces*, *Clostridium* and *Pichia*.

6. The process of claim 5 wherein the microbial host is selected from the group consisting of members of species *E. coli*, *Bacillus subtilis*, *Bacillus licheniformis* and *Pichia pastoris*.

7. The process of claim 6 wherein the microbial host is *E. coli*.

8. The process of claim 1 wherein (a) the transformed microbial host is recombinant *E. coli* DH5.alpha. containing a gene encoding an enzymatically active diol dehydratase enzyme, the gene comprising the DNA sequence of SEQ ID NO. 1; (b) the carbon substrate is glycerol; and (c) the product recovered in step (iii) is 1,3-propanediol.

9. A process for the bioconversion of glycerol to 1,3-propanediol comprising the steps of:

(i) transforming a microbial host selected from the group consisting of the genera *Eschericia*, *Bacillus*, *Klebsiella*, *Citrobacter*, *Saccharomyces*, *Clostridium* and *Pichia* with genes encoding an enzymatically active bacterial diol dehydratase enzyme, the genes derived from a cosmid, the cosmid comprising a DNA fragment of about 35 kb isolated from *Klebsiella pneumoniae*, the cosmid contained within transformed *E. coli* deposited with the American Type Culture Collection under accession number ATCC 69790;

(ii) contacting the transformed microbial host with carbon substrate in a suitable medium; and

(iii) recovering 1,3-propanediol from a suitable medium.

10. The process of claims 8, 1 or 9 wherein the transformed microbial host further contains an alcohol dehydrogenase.

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'CAPLUS' ENTERED AT 09:42:33 ON 17 JUN 1998

COPYRIGHT (C) 1998 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1967 - 17 Jun 1998 VOL 128 ISS 26

FILE LAST UPDATED: 17 Jun 1998 (980617/ED)

FILE 'CAPLUS' ENTERED AT 09:42:33 ON 17 JUN 1998

L1 1240 S REGULON?

L2 2835 S DHA

L3 9 S L1 AND L2

L4 48660 S ANAEROB?

L5 55 S L1 AND L4

L6 16 S L2 AND L4

L7 115385 S FUNG?

L8 16 S L2 AND L7

L9 1604 S L2 NOT (DEHYDROACETIC OR DOCOSAHEX?)(W)ACID

L10 3 S L9 AND L7

L11 94629 S ASPERGILLUS OR SACCHAROMYCES OR YGOSACCHAROMYCES OR PIC

L12 479 S ZYGOSACCHAROMYCES

L13 136506 S DEBARYOMYCES OR MUCOR OR TORULOPSIS OR METHYLOBACTER OR

L14 222716 S L11 OR L12 OR L13

L15 32 S L9 AND L14

L16 219 S 604-63-2P/IT

L17 8 S L14 AND L16

L3 ANSWER 1 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Bioconversion of a fermentable carbon source to 1,3-propanediol by a single microorganism expressing a foreign glycerol or diol dehydratase gene

L3 ANSWER 2 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Phenotypic diversity of anaerobic glycerol dissimilation shown by seven enterobacterial species

L3 ANSWER 3 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Growth temperature-dependent activity of glycerol dehydratase in *Escherichia coli* expressing the *Citrobacter freundii* "dha" "regulon"

L3 ANSWER 4 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Enhancement of 1,3-propanediol production by cofermentation in *Escherichia coli* expressing Klebsiella pneumoniae "dha" "regulon" genes

L3 ANSWER 5 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 1,3-Propanediol production by *Escherichia coli* expressing genes from the Klebsiella pneumoniae "dha" "regulon"

L3 ANSWER 6 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Anaerobic growth of *Escherichia coli* on glycerol by importing genes of the "dha" "regulon" from Klebsiella pneumoniae

L3 ANSWER 7 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Klebsiella pneumoniae 1,3-propanediol NAD<sup>+</sup> oxidoreductase

L3 ANSWER 8 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 Purification and properties of dihydroxyacetone kinase from Klebsiella pneumoniae

L3 ANSWER 9 OF 9 CAPLUS COPYRIGHT 1998 ACS

T1 "dha" System mediating aerobic and anaerobic dissimilation of glycerol in Klebsiella pneumoniae NCIB 418

L5 ANSWER 1 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Mechanism of regulation of 8-hydroxyguanine endonuclease by oxidative stress: roles of FNR, ArcA, and Fur

L5 ANSWER 2 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 The molecular basis for the differential regulation of the hlyE-encoded hemolysin of *Escherichia coli* by FNR and HlyX lies in the improved activating region 1 contact of HlyX

L5 ANSWER 3 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" expression of the *Photobacterium fischeri* lux "regulon" requires the FNR protein which acts upon the left operon

L5 ANSWER 4 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 A promoter and "regulon" specific to the storage organ of sugar beet but that is not regulated by environmental conditions

L5 ANSWER 5 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 HlyX, the FNR homolog of *Acetobacter pleuropneumoniae*, is a [4Fe-4S]-containing oxygen-responsive transcription regulator that "anaerobically" activated FNR-dependent Class I promoters via an enhanced AR1 contact

L5 ANSWER 6 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 The DAN1 gene of *S. cerevisiae* is regulated in parallel with the hypoxic genes, but by a different mechanism

L5 ANSWER 7 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Study of redox-regulated transcription factors in prokaryotes

L5 ANSWER 8 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" expression of the *Vibrio fischeri* lux "regulon" in *E. coli* is FNR-dependent

L5 ANSWER 9 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Methylphosphonic acid degradation and its physiological regulation in *Escherichia coli*

L5 ANSWER 10 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 SoxR, a [2Fe-2S] transcription factor, is active only in its oxidized form

L5 ANSWER 11 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Transcriptional regulation of the *Escherichia coli* rnaT gene

L5 ANSWER 12 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Three two-component signal-transduction systems interact for Pho regulation in *Bacillus subtilis*

L5 ANSWER 13 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Functional significance of the O<sub>2</sub>/ZnSOD in *Escherichia coli*

L5 ANSWER 14 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 The complex bet promoters of *Escherichia coli*: regulation by oxygen (ArcA) choline (BetI), and osmotic stress

L5 ANSWER 15 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Regulators of aerobic and "anaerobic" respiration in *Bacillus subtilis*

L5 ANSWER 16 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 A global signal transduction system regulates aerobic and "anaerobic" CO<sub>2</sub> fixation in *Rhodobacter sphaeroides*

L5 ANSWER 17 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Regulation of "anaerobic" citrate metabolism in Klebsiella pneumoniae

L5 ANSWER 18 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Glutathione is required for maximal transcription of the cobalamin biosynthetic and 1,2-propanediol utilization (cobYpdu) "regulon" and for the catabolism of ethanolamine, 1,2-propanediol, and propionate in *Salmonella typhimurium* LT2

L5 ANSWER 19 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Roles of nitric oxide in inducible resistance of *Escherichia coli* to activated murine macrophages

L5 ANSWER 20 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Aerobic- "anaerobic" gene regulation in *Escherichia coli*: control by the ArcAB and Fnr "regulons"

L5 ANSWER 21 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 The control region of the *ptduscob* "regulon" in *Salmonella typhimurium*

L5 ANSWER 22 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Phenotypic diversity of "anaerobic" glycerol dissimilation shown by seven enterobacterial species

L5 ANSWER 23 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Two global regulatory systems (Cp and Arc) control the cobalamin/propanediol "regulon" of *Salmonella typhimurium*

L5 ANSWER 24 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Control and function of hlyH-RNA synthetases: Diversity and co-ordination

L5 ANSWER 25 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Hyperbaric sensitization of microbes to oxidative stress and disinfection

L5 ANSWER 26 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Choline transport activity in *Staphylococcus aureus* induced by osmotic stress and low phosphate concentrations

L5 ANSWER 27 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Induction of manganese-containing superoxide dismutase in "anaerobic" *Escherichia coli* by diamide and 1,10-phenanthroline: Sites of transcriptional regulation

L5 ANSWER 28 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Expression of extracellular phospholipase from *Serratia liquefaciens* is growth-phase dependent, catabolite-repressed and regulated by "anaerobiosis"

L5 ANSWER 29 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Genetic structure and regulation of the *cysG* gene in *Salmonella typhimurium*

L5 ANSWER 30 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Surface protein-CAT reporter fusions demonstrate differential gene expression in the *vir* "regulon" of *Streptococcus pyogenes*

L5 ANSWER 31 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Regulatory roles of *Fnr*, *Fur*, and *Arc* in expression of manganese-containing superoxide dismutase in *Escherichia coli*

L5 ANSWER 32 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 A single regulatory gene integrates control of vitamin B12 synthesis and propanediol degradation

L5 ANSWER 33 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" induction of the alkylaton-inducible *Escherichia coli* *aidB* gene involves genes of the cysteine biosynthetic pathway

L5 ANSWER 34 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 1,3-Propanediol production by *Escherichia coli* expressing genes from the *Klebsiella pneumoniae* *dha* "regulon"

L5 ANSWER 35 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Adaptation of *Escherichia coli* to respiratory conditions: regulation of gene expression

L5 ANSWER 36 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 A superoxide response "regulon" in *Escherichia coli*

L5 ANSWER 37 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 The *arcB* gene of *Escherichia coli* encodes a sensor-regulator protein for "anaerobic" repression of the *arc* modulon

L5 ANSWER 38 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Substitution of 2 base pairs (1 base pair DNA half-site) within the *Escherichia coli* *lac* promoter DNA site for catabolite gene activator protein places the *lac* promoter in the *FNR* "regulon"

L5 ANSWER 39 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Identification of phosphate starvation-inducible genes in *Escherichia coli* K-12 by DNA sequence analysis of *psi*:*lacZ*(Mu d1) transcriptional fusions

L5 ANSWER 40 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Multiple regulatory elements for the *gla* operon encoding "anaerobic" glycerol-3-phosphate dehydrogenase and the *gpd* operon encoding aerobic glycerol-3-phosphate dehydrogenase in *Escherichia coli*: further characterization of respiratory control

L5 ANSWER 41 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" growth of *Escherichia coli* on glycerol by importing genes of the *dha* "regulon" from *Klebsiella pneumoniae*

L5 ANSWER 42 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 *arcA* (*dye*), a global regulatory gene in *Escherichia coli* mediating repression of enzymes in aerobic pathways

L5 ANSWER 43 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Induction of the manganese-containing superoxide dismutase in *Escherichia coli* is independent of the oxidative stress (*oxyR*-controlled) "regulon"

L5 ANSWER 44 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Transcriptional regulation of *kdsE* in *Escherichia coli* K-12

L5 ANSWER 45 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 A mutant *cap* allele that differentially activates the operons of the *fuc* "regulon" in *Escherichia coli*

L5 ANSWER 46 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Cross-induction of the L-glucose system by L-threonine in *Escherichia coli*

L5 ANSWER 47 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 *Klebsiella pneumoniae* 1,3-propanediol NAD<sup>+</sup>-oxidoreductase

L5 ANSWER 48 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Loss of aldehyde dehydrogenase in an *Escherichia coli* mutant selected for growth on the rare sugar L-galactose

L5 ANSWER 49 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Oxygen regulation in *Salmonella typhimurium*

L5 ANSWER 50 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Overlapping and separate controls on the phosphate "regulon" in *Escherichia coli* K12

L5 ANSWER 51 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 *dha* System mediating aerobic and "anaerobic" dissimilation of glycerol in *Klebsiella pneumoniae* NCIB 418

L5 ANSWER 52 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Synthesis of L-cysteine in *Salmonella typhimurium*

L5 ANSWER 53 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Gene-product relationships of the *nil* "regulon" of *Klebsiella pneumoniae*

L5 ANSWER 54 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 Three kinds of controls affecting the expression of the *gfp* "regulon" in *Escherichia coli*

L5 ANSWER 55 OF 55 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" L<sup>-</sup>alpha-glycerophosphate dehydrogenase of *Escherichia coli*: Its genetic locus and its physiological role

L6 ANSWER 1 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Kinetic study of the oxidation of ascorbic acid by aqueous copper(II) catalyzed by chloride ion

L6 ANSWER 2 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Phenotypic diversity of "anaerobic" glycerol dissimilation shown by seven enterobacterial species

L6 ANSWER 3 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Production of docosahexaenoic acid by marine bacteria isolated from deep sea fish

L6 ANSWER 4 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Specific nutrient transformation processes and change in dehydrogenase activity during formation and evolution of marine detrital microzone

L6 ANSWER 5 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" dihydroxyacetone production from formaldehyde by methanotrophic bacteria

L6 ANSWER 6 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 1,3-Propanediol production by *Escherichia coli* expressing genes from the *Klebsiella pneumoniae* "dha" regulon

L6 ANSWER 7 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Correlations between TTC-dehydrogenase activity and other active parameters during aerobic digestion of excess activated sludge

L6 ANSWER 8 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Microbial enzyme activities: potential use for monitoring decomposition processes

L6 ANSWER 9 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 "Anaerobic" growth of *Escherichia coli* on glycerol by importing genes of the "dha" regulon from *Klebsiella pneumoniae*

L6 ANSWER 10 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Microbial activity measurements for "anaerobic" sludge digestion

L6 ANSWER 11 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 *Klebsiella pneumoniae* 1,3-propanediol NAD<sup>+</sup>-oxidoreductase

L6 ANSWER 12 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Immunochemical properties of NAD<sup>+</sup>-linked glycerol dehydrogenases from *Escherichia coli* and *Klebsiella pneumoniae*

L6 ANSWER 13 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 "dha" System mediating aerobic and "anaerobic" dissimilation of glycerol in *Klebsiella pneumoniae* NCIB 418

L6 ANSWER 14 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Glycerol fermentation in *Klebsiella pneumoniae*: functions of the coenzyme B12-dependent glycerol and diol dehydratases

L6 ANSWER 15 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Effects of storage temperature and duration on total vitamin C content of canned single-strength grapefruit juice

L6 ANSWER 16 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Regulation of glycerol catabolism in *Klebsiella aerogenes*



L6 ANSWER 7 OF 16 CAPLUS COPYRIGHT 1998 ACS

AB During aerobic digestion of excess activated sludge, 2,3,5-triphenyltetrazolium chloride-dehydrogenase activity (TTC-  
"DHA") is significantly correlated to other activity parameters including O uptake rate, microorganism population no., and mixed  
liquor suspended solid concn. . . .

T1 Wastewater treatment (activated-sludge process, excess sludge "anaerobic" digestion in, TTC-dehydrogenase activity  
in)

T1 9082-29-5 RL: PRP (Properties) (activity of, in excess activated sludge "anaerobic" digestion, other activity parameters  
relation to)

L6 ANSWER 15 OF 16 CAPLUS COPYRIGHT 1998 ACS

AB . . . of glycerol as a C source for growth by K. aerogenes strain 2103 involved sep. aerobic (sn-glycerol 3-phosphate  
[G3P]) and "anaerobic" (dihydroxyacetone [""DHA"]) pathways of catabolism. Enzyme and transport activities of the aerobic  
pathway were elevated in cells grown under oxygenated conditions on glycerol or G3P. "Anaerobic" growth on G3P required  
the presence of an exogenous H acceptor such as fumarate; cells thus grown were highly induced in the G3P pathway.  
"Anaerobic" growth on glycerol required no exogenous H acceptors; cells thus grown were highly induced in the "DHA"  
pathway but uninduced in the G3P pathway. The addn. of fumarate electron acceptors did not affect the relative levels of the 2  
pathways. When both glycerol and G3P were provided "anaerobically" with fumarate, the "DHA" pathway was preferentially  
induced, which probably accounts for the exclusive utilization of glycerol until its exhaustion. The presence of a regulatory  
control of the G3P pathway imposed by the operation of the "DHA" pathway was suggested.

T1 Carbon metabolic pathway (for glycerol catabolism, regulation of aerobic and "anaerobic" pathways in Klebsiella  
aerogenes in)

T1 Klebsiella aerogenes (glycerol catabolism by, regulation of aerobic and "anaerobic" pathways in)  
T1 56-81-5, biological studies RL: BPR (Biological process); BIOL (Biological study); PROC (Process) (metab. of, regulation  
of aerobic and "anaerobic" pathways in Klebsiella aerogenes for)

L8 ANSWER 1 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Evaluation of single cell sources of docosahexaenoic acid and arachidonic acid: a 4-week oral safety study in rats

L8 ANSWER 2 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Polyunsaturated fatty acids production by microbial cultivation

L8 ANSWER 3 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Effects of initial sugar concentration and nitrogen sources on the characteristics of growth and fermentation of "fungus" *Thraustochytrium*  
*aureum* ATCC 34304

L8 ANSWER 4 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Molecular cloning, sequence analysis, and functional characterization of the gene *kdsA*, encoding 3-deoxy-D-manno-2- octulosonate-8-  
phosphate synthase of *Chlamydia psittaci* 68C

L8 ANSWER 5 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Plasma fatty acid responses, metabolic effects, and safety of microalgal and "fungal" oils rich in arachidonic and docosahexaenoic acids in  
healthy adults

L8 ANSWER 6 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Improvement of docosahexaenoic acid production in a culture of *Thraustochytrium aureum* by medium optimization

L8 ANSWER 7 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Microbial oils containing arachidonic and docosahexaenoic acids for treating neurological disorders

L8 ANSWER 8 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Dehydroacetic acid and the newly synthesized Schiff base to control aflatoxin accumulation

L8 ANSWER 9 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Concentration of eicosapentaenoic acid and docosahexaenoic acid in an arachidonic acid-producing "fungus", *Moniliella alipina* 1S-4, grown  
with fish oil

L8 ANSWER 10 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Microbial omega-3-containing fats and oils for food use

L8 ANSWER 11 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Lipids of selected molds grown for production of n-3 and n-6 polyunsaturated fatty acids

L8 ANSWER 12 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Production of docosahexaenoic acid by *Thraustochytrium aureum*

L8 ANSWER 13 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Application of a specificity of *Mucor miehei* lipase to concentrate docosahexaenoic acid (""DHA")

L8 ANSWER 14 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Antimycotic activity effect of dehydroacetic acid (""DHA") in food

L8 ANSWER 15 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Side-effects of agrochemicals on soil microorganisms

L8 ANSWER 16 OF 16 CAPLUS COPYRIGHT 1998 ACS

T1 Soft drinks. IV. Preservatives in soft drinks. 3. Transformation of "DHA" [dehydroacetic acid] in citric acid solution on heating and the  
inhibitory effect on *Aspergillus niger*

L10 ANSWER 1 OF 3 CAPLUS COPYRIGHT 1998 ACS

T1 Polyunsaturated fatty acids production by microbial cultivation

L10 ANSWER 2 OF 3 CAPLUS COPYRIGHT 1998 ACS

T1 Molecular cloning, sequence analysis, and functional characterization of the gene *kdsA*, encoding 3-deoxy-D-manno-2- octulosonate-8-  
phosphate synthase of *Chlamydia psittaci* 68C

L10 ANSWER 3 OF 3 CAPLUS COPYRIGHT 1998 ACS

T1 Side-effects of agrochemicals on soil microorganisms

L15 ANSWER 1 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Method for manufacture of eicosapentaenoic acid biver alkyl esters from ester mixtures including hydrolysis with lipase

L15 ANSWER 2 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Composition based on fish oil and containing high levels of polyunsaturated fatty acids and high oxidative stability

L15 ANSWER 3 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Novel transferable .beta.-lactam resistance with cephalosporinase characteristics in "Salmonella" enteritidis

L15 ANSWER 4 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Composting of effluent from a new two-phases centrifuge olive mill. Microbial characterization of the compost

L15 ANSWER 5 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Evidence from whole-sediment, porewater, and elutriate testing in toxicity assessment of contaminated sediments

L15 ANSWER 6 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Molecular cloning, sequence analysis, and functional characterization of the gene *kdsA*, encoding 3-deoxy-D-manno-2- octulosonate-8-  
phosphate synthase of *Chlamydia psittaci* 68C

L15 ANSWER 7 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Bioconversion of a fermentable carbon source to 1,3-propanediol by a single microorganism expressing a foreign glycerol or diol dehydratase  
gene

L15 ANSWER 8 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Biosynthesis of pycholein and dihydroaeruginic acid requires the iron-regulated *pchDCSA* operon in "Pseudomonas" aeruginosa

L15 ANSWER 9 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Bacterial photomutagenicity testing: Distinction between direct, enzyme-mediated and light-induced events

L15 ANSWER 10 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Bioavailability and biodegradation rate of DDT by "Bacillus" sp. B75 in the presence of dissolved humic substances

L15 ANSWER 11 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Biosynthetic pathways of glycerol accumulation under salt stress in "Aspergillus" nidulans

L15 ANSWER 12 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Manufacture of stable and odorless powdered oils and fats

L15 ANSWER 13 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Synthesis of novel phosphatidylhydroxyacetone via transphosphatidyl transfer reaction by phospholipase D

L15 ANSWER 14 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Enrichment of polyunsaturated fatty acids with *Geotrichum candidum* lipase

L15 ANSWER 15 OF 32 CAPLUS COPYRIGHT 1998 ACS

T1 Studies on the inactivation of N-methyl-N'-nitro-N-nitrosoguanidine by the addition of soluble vitamins and SH compounds

- L15 ANSWER 16 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 The antimicrobial effect of a structural variant of subtilin against *Bacillus cereus* T spores and vegetative cells occurs by different mechanisms
- L15 ANSWER 17 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Regulation of glycerol metabolism in *Zygosaccharomyces rouxii* in response to osmotic stress
- L15 ANSWER 18 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Characterization of a glycerol kinase mutant of *Aspergillus niger*
- L15 ANSWER 19 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Classical transketolase functions as the formaldehyde-assimilating enzyme during growth of a dihydroxyacetone synthase-negative mutant of the methylotrophic yeast *Hansenula polymorpha* on mixtures of xylose and methanol in continuous cultures
- L15 ANSWER 20 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Metabolic regulation in the yeast *Hansenula polymorpha*. Growth of dihydroxyacetone kinase/glycerol kinase-negative mutants on mixtures of methanol and xylose in continuous cultures
- L15 ANSWER 21 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Purification and properties of NADP+-dependent glycerol dehydrogenases from *Aspergillus nidulans* and *A. niger*
- L15 ANSWER 22 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Methanol-dependent production of dihydroxyacetone and glycerol by mutants of the methylotrophic yeast *Hansenula polymorpha* blocked in dihydroxyacetone kinase and glycerol kinase
- L15 ANSWER 23 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Genotoxicity of naturally occurring hydroxyanthraquinones
- L15 ANSWER 24 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Dihydroxyacetone kinase from a methylotrophic yeast, *Hansenula polymorpha* CBS 4732. Purification, characterization and physiological role
- L15 ANSWER 25 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Glycerol metabolism in the methylotrophic yeast *Hansenula polymorpha*: phosphorylation as the initial step
- L15 ANSWER 26 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Dihydroxyacetone reductase of a methylotrophic yeast, *Hansenula otuensis*
- L15 ANSWER 27 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Regulation of methanol metabolism in the yeast *Hansenula polymorpha*. Isolation and characterization of mutants blocked in methanol assimilatory enzymes
- L15 ANSWER 28 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Effects of phospholipase C on the  $\beta$ -receptor-adenylate cyclase system of chick erythrocyte membranes
- L15 ANSWER 29 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 In vitro and in vivo studies on the potential mutagenicity of abietane, dihydroxyabietane and abietane epoxide
- L15 ANSWER 30 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 A modified pulse-labelling technique for the detection of early intermediates in microbial metabolism: detection of  $[14C]$ -dihydroxyacetone during assimilation of  $[14C]$ -methanol by *Hansenula polymorpha*
- L15 ANSWER 31 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Intermediates of antibiotics
- L15 ANSWER 32 OF 32 CAPLUS COPYRIGHT 1998 ACS  
T1 Two mutants of glycerol metabolism in *Bacillus subtilis*
- L15 ANSWER 11 OF 32 CAPLUS COPYRIGHT 1998 ACS  
AN 1996:118929 CAPLUS DN 124:170433  
T1 Biosynthetic pathways of glycerol accumulation under salt stress in *Aspergillus nidulans*  
AU Redkar, Rajendra J.; Looy, Robert D.; Singh, Narendra K.  
CS Department of Botany and Microbiology, Auburn University, Auburn, AL 36849-5407, USA  
SO Exp. Mycol. (1995), Volume Date 1995, 19(4), 241-6 CODEN: EXMYD2; ISSN: 0147-5975 DT English  
AB A culture of *Aspergillus nidulans* (FGSC 359) was gradually adapted for growth in media containing up to 2 M NaCl or was exposed to a salt shock with 2 M NaCl. The intracellular glycerol level increased by about 7.9-fold in salt-adapted and 2.4-fold in salt-shocked cultures when compared to the unadapted culture. The biosynthetic pathway involved in the accumulation of glycerol was investigated under long-term salt adaptation and short-term salt shock. Glycerol-3-phosphate dehydrogenase (EC 1.1.1.8) was induced 1.4-fold in salt-shocked but not in salt-adapted cultures. An alternate enzymic pathway involving glycerol
- dehydrogenase (NADP+-dependent) utilizing dihydroxyacetone ( *\*\*\*DHA\** ) and/or DL-glyceraldehyde (DL-GAD) was induced by NaCl. *DHA\**-dependent glycerol dehydrogenase activity was induced about 6.3-fold in salt-adapted and 1.35-fold in salt-shocked cultures, while DL-GAD-dependent activity was induced about 6.1-fold in salt-adapted and 1.2-fold in salt-shocked cultures. However, the level of glycerol dehydrogenase activity with DL-GAD as substrate was 7% of the *DHA\**-dependent activity. We conclude that a salt-inducible NADP+-dependent glycerol dehydrogenase activity electrophoretically indistinguishable from previously described glycerol dehydrogenase I results in glycerol accumulation in salt-stressed *A. nidulans*.
- L15 ANSWER 17 OF 32 CAPLUS COPYRIGHT 1998 ACS  
AN 1992:55338 CAPLUS DN 115:55338  
T1 Regulation of glycerol metabolism in *Zygosaccharomyces rouxii* in response to osmotic stress  
AU Van Zyl, Petrus Jakobus; Prior, Bernard Alexander; Kilian, Stephanus Gouws  
CS Dep. Microbiol. Biochem., Univ. Orange Free State, Bloemfontein, 9300, S. Afr.  
SO Appl. Microbiol. Biotechnol. (1991), 36(3), 369-74 CODEN: AMBIDG; ISSN: 0175-7598 DT English  
AB Enzyme anal. indicated that the metab. of glycerol by *X. rouxii* occurred via either glycerol 3-phosphate (G3P) or dihydroxyacetone ( *DHA\** ). The route via *DHA\** is significant in osmoregulation. The specific activities of glycerol dehydrogenase (GDH-G) and *DHA\** kinase, which metabolize glycerol via *DHA\**, increased 9- and 4-fold, resp., during osmotic stress [0.960 water activity (aw) adjusted with NaCl] when compared to nonstressed conditions (0.998 aw). Both pathways are under metabolic regulation. Glycerol kinase, mitochondrial G3P dehydrogenase, and *DHA\** kinase are induced by glycerol, while the latter is also repressed by glucose. Cells treated with cycloheximide prior to osmotic upshock showed significantly lower *DHA\** kinase and GDH-G levels and lower intracellular glycerol concns. than untreated control cells. Thus protein synthesis is essential for osmotic adaptation.
- L15 ANSWER 18 OF 32 CAPLUS COPYRIGHT 1998 ACS  
AN 1991:38939 CAPLUS DN 114:38939  
T1 Characterization of a glycerol kinase mutant of *Aspergillus niger*  
AU Witteveen, Cor F. B.; Van de Vondervoort, Peter; Dijkema, Cor; Swart, Klaas; Visser, Jaap  
CS Dep. Genet., Agric. Univ., Wageningen, 6703 HA, Neth.  
LA English  
AB A glycerol-kinase-deficient mutant of *A. niger* was isolated. Genetic anal. revealed that the mutation is located on linkage group VI. The phenotype of this mutant differed from that of a glycerol kinase mutant of *A. nidulans* in its ability to utilize dihydroxyacetone ( *\*\*\*DHA\** ). The weak growth on glycerol of the *A. niger* glycerol kinase mutant showed that glycerol phosphorylation is an important step in glycerol catabolism. The mutant could still grow normally on *DHA\** because of the presence of a *DHA\** kinase. This enzyme, probably in combination with an NAD+-dependent glycerol dehydrogenase, present only in the mutant, is responsible for the weak growth of the mutant on glycerol. Enzymic anal. of both the mutant and the parental strain showed that glucose 3 different glycerol dehydrogenases were formed under different physiol. conditions: the NAD+-dependent enzyme described above, a constitutive NADP+-dependent enzyme, and a D-glyceraldehyde-specific enzyme induced on D-galacturonate. The glycerol kinase mutant showed impaired growth on D-galacturonate.
- L15 ANSWER 19 OF 32 CAPLUS COPYRIGHT 1998 ACS  
AN 1990:474497 CAPLUS DN 113:74497  
T1 Classical transketolase functions as the formaldehyde-assimilating enzyme during growth of a dihydroxyacetone synthase-negative mutant of the methylotrophic yeast *Hansenula polymorpha* on mixtures of xylose and methanol in continuous cultures  
AU De Koning, W.; Bonting, K.; Harder, W.; Dijkhuizen, L.  
CS Dep. Microbiol., Univ. Groningen, Haren, 9751 NN, Neth.  
SO Yeast (1990), 6(2), 117-25 CODEN: YESTE3; ISSN: 0749-503X DT English  
AB Contrary to expectation, a mutant of *H. polymorpha* blocked in dihydroxyacetone ( *\*\*\*DHA\** ) synthase was able to assimilate methanol-carbon when grown in chemostat culture on mixts. of xylose and methanol. Incubation of a *DHA\**-synthase- and *DHA\** kinase-neg. double mutant resulted in *DHA\** accumulation, indicating that a *DHA\** synthase-type of reaction was involved. Low residual *DHA\** synthase activity subsequently was present when using an assay with improved sensitivity. This activity was not assoc. with the (mutated) *DHA\** synthase protein, which was still present in the peroxisomes, but with the enzyme transketolase. Transketolase from methanol-grown cells was purified [525-fold] to homogeneity in 9% yield. The native enzyme was dimeric, as has been reported for other transketolases, with a subunit mol. wt. of 74,000. The affinity of the purified enzyme for formaldehyde was low ( $K_m = 5$  mM), but high for xylulose-5-phosphate (10  $\mu$ M). The *in vivo* functioning of transketolase in formaldehyde assimilation, and the influence of the hydration state of formaldehyde is discussed.
- L15 ANSWER 20 OF 32 CAPLUS COPYRIGHT 1998 ACS  
AN 1990:474496 CAPLUS DN 113:74496  
T1 Metabolic regulation in the yeast *Hansenula polymorpha*. Growth of dihydroxyacetone kinase/glycerol kinase-negative mutants on mixtures of methanol and xylose in continuous cultures  
AU De Koning, W.; Weusthuis, R. A.; Harder, W.; Dijkhuizen, L.

CS Dep. Microbiol., Univ. Groningen, Haren, 9751 NN, Neth.

SO Yeast (1990), 6(2), 107-115 CODEN: YESTEJ ISSN: 0749-503X DT Journal LA English

AB The physiological responses of H. polymorpha wild-type and mutant strains 17B (dihydroxyacetone kinase-neg.) and 17BC51 (dihydroxyacetone kinase- and glycerol kinase-neg.) to growth on mixts. of xylose and methanol in chemostats were investigated. Increasing methanol concns. (0-110 mM) in the feed of the wild-type culture resulted in increasing cell densities and a gradual switch towards methanol metab. At the lower methanol feed concns. the mutant cultures used methanol and xylose to completion and changes in enzyme patterns comparable to the wild type were obsd. This was not reflected in significant changes in cell densities. Instead, formaldehyde assimilation resulted in dihydroxyacetone ( "DHA" ) prodn., which was proportional to the amt. of methanol added. At intermediate methanol concns., the culture showed a strong variation in "DHA" levels and cell densities. Further increases in the methanol feed concns. resulted in a drop in "DHA" accumulation rates, repression of alc. oxidase synthess, and accumulation of residual methanol. These phenomena were studied in more detail in transition expts. and with gradients of methanol. The results indicate that xylose-5-phosphate (Xu5P) generated in xylose metab. served as acceptor mol. for formaldehyde assimilation by the peroxisomal enzyme "DHA" synthase. Accumulation of "DHA" in the mutant cultures, however, further diminished the availability of carbon for growth. Thus, with increasing methanol concns., Xu5P eventually became growth rate-limiting. This resulted in an unstable situation but wash-out of the culture did not occur to a significant extent. Instead, "DHA" accumulation ceases and cell densities, and the enzymes specifically involved in xylose metab. increase, indicating that the organism resumed its xylose metab. The mol. mechanism controlling the partitioning of Xu5P over xylose (pentose phosphate pathway) and methanol (peroxisome) metab. under these conditions remain to be elucidated.

L15 ANSWER 21 OF 32 CAPLUS COPYRIGHT 1988 ACS

AN 1990:454844 CAPLUS DN 113:54844

T1 Purification and properties of NAD(P)-dependent glycerol dehydrogenases from "Aspergillus" nidulans and A. niger

AU Schuurink, R.; Busink, R.; Hondmann, D. H. A.; Witteveen, C. F. B.; Visser, J.

CS Dep. Geneet., Agric. Univ., Wageningen, 6703 HA, Neth.

SO J Gen. Microbiol. (1990), 136(6), 1043-50 CODEN: JGMIAN; ISSN: 0022-1287 DT Journal LA English

AB Glycerol dehydrogenase (NADP-specific; EC 1.1.1.72) (I) was purified from mycelium of A. nidulans and A. niger using different purifn. procedures. Both enzymes had a mol. wt. of approx. 38,000 and were immunol. cross-reactive, but had different amino acid compns. and pI values. For both enzymes, the substrate specificity was limited to glycerol and erythritol for the oxidative reaction and to dihydroxyacetone ( "DHA" ), diacetyl, methylglyoxal, erythrose, and D-glyceraldehyde for the reductive reaction. I of A. nidulans had a turnover no. twice that of A. niger I at pH 6.0, whereas inhibition by NADP was less (Ki = 45 vs. 13 .mu.M). It was proposed that both enzymes catalyze in vivo the reductn. of "DHA" to glycerol and that they are regulated by the anabolic reductn. charge.

L15 ANSWER 22 OF 32 CAPLUS COPYRIGHT 1988 ACS

AN 1990:215157 CAPLUS DN 112:215157

T1 Methanol-dependent production of dihydroxyacetone and glycerol by mutants of the methylotrophic yeast "Hansenula" polymorpha blocked in dihydroxyacetone kinase and glycerol kinase

AU De Koning, W.; Weuhsuijs, R. A.; Harder, W.; Dijkhuizen, L.

CS Dep. Microbiol., Univ. Groningen, Haren, NL-9751 NN, Neth.

SO Appl. Microbiol. Biotechnol. (1990), 32(6), 693-8 CODEN: AMBIDG; ISSN: 0175-7598 DT Journal LA English

AB Various factors controlling dihydroxyacetone ( "DHA" ) and glycerol prodn. from methanol by resting cell suspensions of a mutant of H. polymorpha blocked in "DHA" kinase and glycerol kinase, were investigated. The presence of methanol (250 mM) and an addnl. substrate (0.5%, w/v) to replenish the xylose-5-phosphate required for the assimilation reaction (DHQa synthase) was essential for significant triose prodn. by this double mutant. A no. of sugars were tested as addnl. substrates and C5 sugars gave the highest triose accumulation (ca. 20 mM after 45 h). Glucose was the poorest addnl. substrate and triose prodn. only started after its exhaustion, which occurred in the first few hours. Other sugars were metabolized at a much lower rate and accumulation of trioses began right at the start of the expts. and gradually increased with time. The prodn. rate of total trioses increased, and the relative amt. of glycerol diminished with higher oxygen supply rates. The data suggest that conversion of "DHA" into glycerol, catalyzed by reduced nicotinic adenine dinucleotide (NADH)-dependent "DHA" reductase, is partly regulated via intracellular NADH levels. Further support for this hypothesis was obtained in expts. with antimycin A, an inhibitor of the electron transport chain. Addn. of higher amts. of methanol and xylose, either by increasing the initial concns. or by repeated addn. of these substrates, resulted in considerably enhanced productivity and a switch towards glycerol formation. After reaching a level of approx. 25 mM the "DHA" concn. remained const. while the glycerol level gradually increased with time. After an incubation period of 350 h, a total of 3.9 M methanol and 0.62 M xylose had been converted, which resulted in accumulation of 0.76 M trioses, mostly glycerol.

L15 ANSWER 25 OF 32 CAPLUS COPYRIGHT 1988 ACS

AN 1987:614645 CAPLUS DN 107:214645

T1 Glycerol metabolism in the methylotrophic yeast "Hansenula" polymorpha: phosphorylation as the initial step

AU De Koning, W.; Harder, W.; Dijkhuizen, L.

CS Dep. Microbiol., Univ. Groningen, Haren, NL-9751 NN, Neth.

SO Arch. Microbiol. (1987), 148(4), 314-20 CODEN: AMICOW; ISSN: 0302-8933 DT Journal LA English

AB In H. polymorpha glycerol is metabolized via glycerol kinase and NAD(P)-independent glycerol 3-phosphate (G3P) dehydrogenase, enzymes which hitherto were reported to be absent in this methylotrophic yeast. Activity of glycerol kinase was readily detectable when cell-free exts. were incubated at pH 7.8 with glycerol, ATP, and Mg2+ and a discontinuous assay for G3P formation was used. This glycerol kinase activity could be sepd. from dihydroxyacetone ( "DHA" ) kinase activity by ion exchange chromatog. Glycerol kinase showed relatively low affinities for glycerol (apparent Km = 1.0 mM) and ATP (apparent Km = 0.5 mM) and was not active with other substrates tested. No inhibition by fructose 1,6-bisphosphate (FBP) was obsd. Both NAD-dependent and NAD(P)-independent G3P dehydrogenases were present. Glucose partly repressed synthesis of glycerol kinase and NAD(P)-independent G3P dehydrogenase, but compared to several other non-repressing C sources no clear induction of these enzymes by glycerol was apparent. Among glycerol-neg. mutants of H. polymorpha strain 17B (a "DHA" kinase-neg. mutant), strains blocked in either glycerol kinase or membrane-bound G3P dehydrogenase were identified. Crosses between representatives of the latter mutants and wild type resulted in the isolation of, among others, segregants which had regained "DHA" kinase but were still blocked in the membrane-bound G3P dehydrogenase. These strains, employing the oxidative pathway, were only able to grow very slowly in glycerol mineral medium.

L15 ANSWER 26 OF 32 CAPLUS COPYRIGHT 1988 ACS

AN 1987:613959 CAPLUS DN 107:213959

T1 Dihydroxyacetone reductase of a methylotrophic yeast, "Hansenula" otaunensis

AU Yamada, Keiko; Tani, Yoshiki

CS Fac. Agric., Kyoto Univ., Kyoto, 606, Japan

SO Agric. Biol. Chem. (1987), 51(9), 2629-31 CODEN: ABCH46; ISSN: 0002-1369 DT Journal LA English

AB Dihydroxyacetone ( "DHA" ) reductase formation was induced by growing H. otaunensis in MeOH-contg. media, and enzyme-substrate activity was tested. "DHA" reductase was isolated by applying cell-free ext. to a DEAE-cellulose column and eluting with KCl in buffer. Oxidative activity, in the presence of NAD+, was low toward glycerol (Km = 2.9 mM) and higher toward 1,2-propanediol and EtOH. The enzyme had reductive activity, in the presence of NADH+, toward "DHA" (Km = 0.36 mM), methylglyoxal, and acetal.

L15 ANSWER 32 OF 32 CAPLUS COPYRIGHT 1988 ACS

AN 1972:472470 CAPLUS DN 77:72470

T1 Two mutants of glycerol metabolism in "Bacillus" subtilis

AU Sateb, S. A.

CS Serv. Physiol. Cellulaire, Inst. Pasteur, Paris, Fr.

SO Can. J. Microbiol. (1972), 18(6), 1315-25 CODEN: CJMAZ DT Journal LA French

AB Two pathways for the degradation of glycerol are found in B. subtilis 168. Each pathway includes two enzymes which can catalyze the formation of dihydroxyacetone phosphate from glycerol in vitro. The first pathway includes a glycerol dehydrogenase (gl-D) and a dihydroxyacetone kinase ( "dhk" -K). The second pathway includes a glycerol kinase (gl-K) and an alpha.-glycerophosphate dehydrogenase (glp-D). Enzymes of both pathways are repressed in the presence of glucose. Only the enzymes of the second pathway are inducible. The inducer is probably glycerophosphate, utilization of which as a C source by B. subtilis is demonstrated. Degradation of glycerol in B. subtilis proceeds through the second pathway. This was demonstrated by the isolation of a mutant (gl-2) impaired in glycerol kinase, and which cannot use glycerol as a C source. Another mutant (gl-1) was isolated, which cannot use glycerol as a C source. When comparing the activity of the four enzymatically particularly gl-K, no significant differences were observed between the wild strain and the mutant gl-1. This indicates the existence of a glycerol permeation system in B. subtilis. A mutation affecting this system would explain the behavior of the mutant gl-1.

L17 ANSWER 1 OF 8 CAPLUS COPYRIGHT 1988 ACS

T1 Metabolic engineering of propanediol pathways

L17 ANSWER 2 OF 8 CAPLUS COPYRIGHT 1988 ACS

T1 Metabolic engineering of an improved 1,3-propanediol fermentation ("Klebsiella pneumoniae", "Bacillus" licheniformis)

L17 ANSWER 3 OF 8 CAPLUS COPYRIGHT 1988 ACS

T1 Production of 1,3-propanediol from glycerol by recombinant bacteria expressing recombinant diol dehydratase

L17 ANSWER 4 OF 8 CAPLUS COPYRIGHT 1988 ACS

T1 Bioconversion of a fermentable carbon source to 1,3-propanediol by a single microorganism expressing a foreign glycerol diol dehydratase gene

L17 ANSWER 5 OF 8 CAPLUS COPYRIGHT 1988 ACS

T1 Process for making 1,3-propanediol from carbohydrates using mixed microbial cultures

L17 ANSWER 6 OF 8 CAPLUS COPYRIGHT 1988 ACS

AN 1993:146193 CAPLUS DN 118:146193

TI Microbial production and downstream processing of 2,3-butanediol  
 AU Altschlar, A. S.; Vaz, Russell, C. E.; Jonas, R.; Crantio, A. Quesada; Schaller, K.  
 CS GBF-Ges Biotechnol. Forsch. mBH Braunschweig, W-3300, Germany  
 SO J. Biotechnol. (1993), 27(3), 317-29 CODEN: JBTID4; ISSN: 0168-1656 DT Journal LA English  
 AB In the direct conversion of starch by "Bacillus" polymyxa a max. of 38 g 2,3-butanediol/L is produced, with a yield of 0.28 g  
 diol/g starch. By preliminary saccharification of starch and then cultivation with Klebsiella oxytoca, a 2,3-butanediol concn. of  
 99-100 g/L is achieved with a yield of 0.5 g diol/g starch. K. oxytoca converts high-test molasses to 2,3-butanediol in the same  
 concn. and yield. The same diol concn., only at lower productivity, can also be achieved by conversion of black strap molasses,  
 provided it contains <2% salts. 2,3-Butanediol can be sepd. from bioprocess media with very good results by salting out using  
 anhyd. K2CO3. After preheating the medium from molasses or saccharified starch conversion process, it was possible to sep.  
 94-96% of the 2,3-butanediol using 53-56% K2CO3. The concn. of the 2,3-butanediol in the resulting diol phase was 97%.  
 Sailing out can also be used to sep. other diols produced using microbial methods.

L17 ANSWER 7 OF 8 CAPLUS COPYRIGHT 1998 ACS  
 AN 1990.234037 CAPLUS DN 112.234037  
 TI Fermentative manufacture of 1,3-propanediol from glycerol  
 PA Kretschmann, Josef; Carduck, Franz Josef; Deckwer, Wolf Dieter; Tag, Carmen  
 SO Ger. Offen., 7 pp. CODEN: GWXXBX  
 PI DE 3829618 A1 900315 AI DE 88.3829618 880901 DT Patent LA German  
 AB Propane-1,3-diol is manufd. from a glycerol-contg. soln. (5-20% by wt.) with a microorganism such as Clostridium,  
 Enterobacterium, Lactobacillus, "Bacillus", Citrobacter, or Klebsiella in a yield of g/req. 0.5 g/hL. Klebsiella pneumoniae DSM  
 2026 was batch-cultured at 37 degree, under anaerobic conditions to yield a max. of 2.3 g propane-1,3-diol from a starting  
 glycerol concn. of 100 g/L; other glycerol concns. (50-200 g/L) produced lower yields.

L17 ANSWER 8 OF 8 CAPLUS COPYRIGHT 1998 ACS  
 AN 1983.214106 CAPLUS DN 98.214106  
 TI Neutral solvent production from halophilic, photolithotrophically grown algae by linked fermentations  
 AU Nakas, J. P.; Schaefer, M.; Parkinson, C. M.; Conley, C. E.; Tanenbaum, S. W.  
 CS Coll. Environ. Sci. For., SUNY, Syracuse, NY, 13210, USA  
 SO Comm. Eur. Communities, [Rep.] EUR (1983), EUR 8245, Energy Biomass, 298-302 CODEN: CECE99 DT Report LA  
 English  
 AB Five species of Dunaliella were examd. for glycerol [56-81.5] accumulation, growth rate, cell d., and protein and chlorophyll  
 content. The suitability of each algal species for such bioconversions was judged according to glycerol accumulation and  
 quantities of neutral solvents produced after sequential bacterial ferms. When grown in 2M NaCl, with 24 mM NaHCO3 or 3%  
 CO2 at 28 degree, and with 25,000 lx at container surface, 4 of the 5 species tested (D. tertiolecta, D. parva, and  
 D. bardawil) produced 10-20 mg of glycerol/L. A Clostridium converted an algal biomass mixt. supplemented with 4% glycerol  
 to approx. 18 g/L of mixed alcs. (EtOH [64-17.5], 1,3-propanediol [504-63-2], and BuOH [71-36-3]). Acetone was not detected.  
 A soil isolate, tentatively classified as a member of the genus "Bacillus", converts glycerol into EtOH at a final concn. of 7.0-9.6  
 g/L. An enrichment culture from sewage sludge resolved to contain 2 gram-neg. rods converts the algal biomass-glycerol mixt.  
 solely to 1,3-propanediol [504-63-2] at a final concn. of 4.2-5.3 g/L. Addnl., Dunaliella concs. of ltoreq. 200-fold, can be directly  
 fermented to mixed solvents.

..... STN Columbus .....  
 FILE 'BIOSIS' ENTERED AT 09:30:06 ON 22 JUN 1998 COPYRIGHT (C) 1998 BIOSIS(R)  
 FILE COVERS 1969 TO DATE.  
 CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNS) PRESENT FROM JANUARY 1969 TO  
 DATE.

FILE 'MEDLINE' ENTERED AT 09:32:26 ON 22 JUN 1998  
 FILE LAST UPDATED: 18 JUN 1998 (19980618/UP). FILE COVERS 1966 TO DATE.

FILE 'BIOSIS' ENTERED AT 09:30:06 ON 22 JUN 1998  
 L1 45281 S SALMONELLA  
 L2 246 S SENFTENBERG  
 L3 236 S L1 AND L2  
 L4 28504 S GLYCEROL  
 L5 3679 S GLYCERIN  
 L6 1 S (L4 OR L5) AND L3  
 L7 331 S DERBY  
 L8 0 S L7 AND (L4 OR L5) AND L1  
 FILE 'MEDLINE' ENTERED AT 09:32:26 ON 22 JUN 1998

L9 3 S L6  
 L10 0 S L8  
 L11 85 S L3  
 L12 280 S ENTERICA  
 L13 1 L12 AND L11  
 L14 0 L7 AND L12  
 L15 68 L7 AND L1

L6 ANSWER 1 OF 1 BIOSIS COPYRIGHT 1998 BIOSIS  
 AN 93.96603 BIOSIS  
 DN BA95.51799  
 TI NOVOBIOCIN BRILLANT GREEN "GLYCEROL" LACTOSE AGAR FURTHER  
 ROUTINE EVALUATION ON 5554 HUMAN STOOLS AND 982 VETERINARY SAMPLES.  
 AU POISSON D.M.; NUGIER J.P.; FLORENCE S.; BELLAOUNI H  
 CS LAB. BACTERIOL. CHRO, BP 2439, 45032 ORLEANS CEDEX, FRANCE.  
 SO PATHOL. BIOL. 40 (8), 1992, 793-796. CODEN: PTBIAN ISSN: 0031-3009  
 LA English  
 AB In order to provide a wider evaluation of "Novobiocin-brilliant green-glycerol" lactose (NBGL) agar, dishes of this medium  
 were added to standard media: Hektoen (H), "Salmonella"-Shigella agar (SS), at all plating steps for 5554 stool cultures of  
 human medical routine (280 isolates) and 982 samples of veterinary routine (133 isolates). NBGL expectively missed lactose  
 \*\*\*glycerol\*\*\* positive strains of the serotype "Senftenberg" (n = 4), H2S negative strains (n = 1), and strains of the Typhi  
 serotype (n = 7). Otherwise, three strains, of serotype Virchow, were unable to grow on NBGL (0.7% of positive samples).  
 Nevertheless overall sensitivities were increased by approx. 10% in the human routine (H: 70%; SS: 63%; NBGL: 94%; at the  
 direct plating step) (H: 83%; SS: 84%; NBGL: 92%; at the enrichment plating step) and by 48% in the veterinary one (NBGL:  
 97%; versus usual media: 68%). Positive predictive values of black centered colonies were significantly higher on NBGL in  
 human routine (H: 38%; SS: 40%; NBGL: 85%; at the direct plating step) (H: 20%; SS: 21%; NBGL: 82%; at the enrichment  
 plating step); and in the veterinary one as well (NBGL: 90%; versus usual media: 17%). These data suggest that NBGL agar  
 does improve "Salmonella" isolation in these kinds of routines, and that growth should be made sure before experiments using  
 given strains.  
 ST METHOD  
 RN 56-81-5 (GLYCEROL) 63-42-3 (LACTOSE) 303-81-1 (NOVOBIOCIN)  
 9002-18-0 (AGAR)  
 CC Biochemical Studies-Carbohydrates 10068 Pharmacology-General \*22002 Microbiological Apparatus, Methods and Media \*32000  
 Veterinary Science-Microbiology \*38006 Chemotherapy-Antibacterial Agents \*38504  
 BC Animals-Unspecified 33000 Horminides 86215

L9 ANSWER 1 OF 3 MEDLINE AN 95130176 MEDLINE DN 95130176  
 TI Development of a conjoint phage typing & biotyping schema for "Salmonella" enterica serovar "Senftenberg" (S.  
 "senftenberg") & the correlation of biotypes with phage types.  
 AU Kumar S, Sharma N C; Singh S, Bhatia R, Singh H  
 CS Department of Medical Microbiology, Postgraduate Institute of Medical Education & Research, Chandigarh.  
 SO INDIAN JOURNAL OF MEDICAL RESEARCH, (1994 Dec) 100 257-61. Journal code: G.J.F. ISSN: 0971-5916. CY  
 DT Journal Article; (JOURNAL ARTICLE) LA English EM 199504  
 AB A total of 287 strains of S. "senftenberg" received from various parts of India during 1969 to 1992 were phage typed using  
 six lysogenic phages. The typability was 90.3 per cent and 14 different phage types could be defined excluding a small group of  
 untypable strains. A biotyping scheme was developed utilising six characters and 13 biotypes could be defined. Stern's  
 "glycerol" medium proved to be the best discriminatory medium. Diversity indexes of phage typing and biotyping schemes were  
 0.868 and 0.503 respectively. Better discrimination was obtained when phage types were subdivided into different biotypes with  
 a diversity index of 0.931. The schemes were found stable, reproducible and epidemiologically useful.  
 CT \*Bacterial Typing Techniques \*Bacteriophage Typing Lysogeny \*\*Salmonella: Q, classification \* Salmonella: VI, virology \* Salmonella  
 Phages: PH, physiology

L9 ANSWER 3 OF 3 MEDLINE AN 92160256 MEDLINE DN 92160256  
 TI Differentiation of "Salmonella" "senftenberg" into biogroups.  
 AU Tsuchi L M; McLaren I M; Smith J E; Wray C  
 CS Central Veterinary Research Institute, Lusaka, Zambia.  
 SO VETERINARY RECORD, (1991 Dec 14) 129 (24) 530-1. Journal code: XBS. ISSN: 0042-4900. CY ENGLAND: United  
 Kingdom  
 DT Journal Article; (JOURNAL ARTICLE) LA English FS Priority Journals EM 199205  
 AB Ninety-six strains of "Salmonella" "senftenberg", isolated between 1984 and 1986 from different parts of England and  
 Wales, were tested for their biochemical reactions and biotyped according to the method of Duguid and others (1975). Nine  
 biogroups were identified on the basis of their metabolism of L-tartrate, D-tartrate, Bitter's xylose and Stern's "glycerol". In

addition, fumaric, oxalic, succinic, glutaric, malonic, maleic, L-malic, L-aspartic, lactic and formic acids were used but did not increase the discrimination. Three biogroups (7, 2 and 5) accounted for 79 per cent of the cultures examined.

L11 ANSWER 1 OF 85 MEDLINE  
T1 National outbreak of "Salmonella" "senftenberg" associated with infant food.

L11 ANSWER 2 OF 85 MEDLINE  
T1 A plasmid-mediated CMY-2 beta-lactamase from an Algerian clinical isolate of "Salmonella" "senftenberg".

L11 ANSWER 3 OF 85 MEDLINE  
T1 CT18cr94, a conjugative transposon found in enterobacteria.

L11 ANSWER 4 OF 85 MEDLINE  
T1 Naturally occurring deletions in the centisome 63 pathogenicity island of environmental isolates of "Salmonella" spp.

L11 ANSWER 5 OF 85 MEDLINE  
T1 Isolation of "Salmonella" "senftenberg" bacteriophages.

L11 ANSWER 6 OF 85 MEDLINE  
T1 Characteristics of "Salmonella" strains isolated from sporadic diarrheal cases during 1992-1994 in the Philippines.

L11 ANSWER 7 OF 85 MEDLINE  
T1 Nosocomial outbreak of gastroenteritis due to "Salmonella" "senftenberg".

L11 ANSWER 8 OF 85 MEDLINE  
T1 In vitro fructooligosaccharide utilization and inhibition of "Salmonella" spp. by selected bacteria.

L11 ANSWER 9 OF 85 MEDLINE  
T1 Transmission of lethal "Salmonella" "senftenberg" from mother's breast-milk to her baby.

L11 ANSWER 10 OF 85 MEDLINE  
T1 Opreloxacin resistance among multidrug resistant strains of "Salmonella" "senftenberg" in India [letter].

L11 ANSWER 11 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg": epidemics in India and present status.

L11 ANSWER 12 OF 85 MEDLINE  
T1 "Salmonella" group-E ( "senftenberg" ) lung abscess: a case report.

L11 ANSWER 13 OF 85 MEDLINE  
T1 Development of a conjugate phage typing & biotyping scheme for "Salmonella" enterica serovar "Senftenberg" (S. "senftenberg" ) & the correlation of biotypes with phage types.

L11 ANSWER 14 OF 85 MEDLINE  
T1 Correlation of phospholipase A & enterotoxin production by "Salmonella" typhimurium with reference to virulence parameters.

L11 ANSWER 15 OF 85 MEDLINE  
T1 Survival of "Salmonella" "senftenberg" and "Salmonella" typhimurium in glassy and rubbery states of gelatin.

L11 ANSWER 16 OF 85 MEDLINE  
T1 Survival of *Salmonellas* in composted and not composted solid animal manure.

L11 ANSWER 17 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg" septicemia: a nursery outbreak.

L11 ANSWER 18 OF 85 MEDLINE  
T1 Serotypes of "Salmonella" isolated from California turkey flocks and their environment in 1984-89 and comparison with human isolates.

L11 ANSWER 19 OF 85 MEDLINE  
T1 The occurrence of salmonellae in bean sprouts in Thailand.

L11 ANSWER 20 OF 85 MEDLINE  
T1 Transferable drug resistance in "Salmonella" "senftenberg".

L11 ANSWER 21 OF 85 MEDLINE  
T1 Incidence of "salmonella" meningitis in Ludhiana (Punjab).

L11 ANSWER 22 OF 85 MEDLINE  
T1 "Salmonella"-induced enteritis. Clinical, serotypes and treatment.

L11 ANSWER 23 OF 85 MEDLINE  
T1 Asymptomatic "Salmonella" "senftenberg" carriage in a neonatal ward.

L11 ANSWER 24 OF 85 MEDLINE  
T1 Use of ribotyping for characterization of "Salmonella" serotypes.

L11 ANSWER 25 OF 85 MEDLINE  
T1 Novobiocin-brilliant green-glycerol-lactose-agar: further routine evaluation on 5554 human stools and 982 veterinary samples.

L11 ANSWER 26 OF 85 MEDLINE  
T1 Amplification of an invA gene sequence of "Salmonella" typhimurium by polymerase chain reaction as a specific method of detection of "Salmonella".

L11 ANSWER 27 OF 85 MEDLINE  
T1 "Salmonella" enteritidis-specific monoclonal antibodies.

L11 ANSWER 28 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg" carrier state in a neonate following septicemia [letter].

L11 ANSWER 29 OF 85 MEDLINE  
T1 [Kinetics of the inactivation of "Salmonella" during thermal disinfection of liquid manure]. Kinetik der Inaktivierung von Salmonellen bei der thermischen Desinfektion von Flüssigmist.

L11 ANSWER 30 OF 85 MEDLINE  
T1 Differentiation of "Salmonella" "senftenberg" into biogroups.

L11 ANSWER 31 OF 85 MEDLINE  
T1 Isolation of "Salmonella" "senftenberg" from different clinical sources.

L11 ANSWER 32 OF 85 MEDLINE  
T1 Nosocomial infection due to "salmonella" "senftenberg" (case report).

L11 ANSWER 33 OF 85 MEDLINE  
T1 Decreased "Salmonella" colonization in turkey poultis inoculated with anaerobic cecal microflora and provided dietary lactose.

L11 ANSWER 34 OF 85 MEDLINE  
T1 [Disinfection studies with "Salmonella" "senftenberg" using egg shells as germ carriers]. Desinfektionsversuche mit "Salmonella" "senftenberg" unter Verwendung von Eischalen als Keimträger.

L11 ANSWER 35 OF 85 MEDLINE  
T1 Extent of salmonellae contamination in breeder hatcheries.

L11 ANSWER 36 OF 85 MEDLINE  
T1 Development and application of an ELISA for detecting antibodies to "Salmonella" enteritidis in chicken flocks.

L11 ANSWER 37 OF 85 MEDLINE  
T1 Production and characterization of a monoclonal antibody specific for "Salmonella" O19-antigen.

L11 ANSWER 38 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg" outbreak in a neonatal unit.

L11 ANSWER 39 OF 85 MEDLINE  
T1 Sandwich enzyme immunoassays for detection of "Salmonella" typhi.

L11 ANSWER 40 OF 85 MEDLINE  
T1 A comparative study of the heat resistance of salmonellas in homogenized whole egg, egg yolk or albumen.

L11 ANSWER 41 OF 85 MEDLINE  
T1 Osteomyelitis: a rare complications of "Salmonella" "senftenberg" infection—a case report.

L11 ANSWER 42 OF 85 MEDLINE  
T1 Evaluation of coagulination test for serotyping of enteropathogenic bacteria.

L11 ANSWER 43 OF 85 MEDLINE  
T1 The survival of salmonellas in shell eggs cooked under simulated domestic conditions.

L11 ANSWER 44 OF 85 MEDLINE  
T1 Effect of a new pelleting process on the level of contamination of poultry mash by *Escherichia coli* and "Salmonella".

L11 ANSWER 45 OF 85 MEDLINE

T1 [The tenacity of bacteria in the airborne state. V1. Tenacity of airborne S. "senftenberg" ]. Die Tenazität von Bakterien im Luftgetragenen Zustand. VI. Mitteilung: Tenazität Luftgetragener S. "senftenberg" .

L11 ANSWER 46 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg" epidemic in a neonatal nursery.

L11 ANSWER 47 OF 85 MEDLINE  
T1 [Species structure of Salmonellas isolated from mammals, poultry, feed mixtures and the environment 1976-1980]. Vidova struktura na salmonelle, izolirani od bozaintisi, ptici, furežni smesi i vrinista sreda za perioda 1976-1980 g.

L11 ANSWER 48 OF 85 MEDLINE  
T1 [Sensitivity of Salmonellas isolated from poultry to bacteriophage O1]. Otvorljivost na salmonela, izolirani od ptici, kum bakteriofag O1.

L11 ANSWER 49 OF 85 MEDLINE  
T1 "Salmonella" shed by horses with colic.

L11 ANSWER 50 OF 85 MEDLINE  
T1 Molecular relationships between virulence plasmids of "Salmonella" serotypes typhimurium and dublin and large plasmids of other "Salmonella" serotypes.

L11 ANSWER 51 OF 85 MEDLINE  
T1 [Survival of salmonellas and ascaris eggs during sludge utilization in forestry (author's trans)]. Untersuchungen über die Tenazität von Salmonellen und Ascaridenreiem bei der Ausbringung von Klärschlamm in Waldbeständen.

L11 ANSWER 52 OF 85 MEDLINE  
T1 Salt extends the upper temperature limit for growth of food-poisoning bacteria.

L11 ANSWER 53 OF 85 MEDLINE  
T1 Behavior of pathogenic bacteria in the oyster, Crassostrea commercialis, during depuration, re-baying, and storage.

L11 ANSWER 54 OF 85 MEDLINE  
T1 [ ""Salmonella" destruction by heating during the customary preparation of dehydrated food products (author's trans)]. "Salmonella" - Abtötung durch Wärme bei der Zubereitung von Lebensmitteln/Trockenprodukten.

L11 ANSWER 55 OF 85 MEDLINE  
T1 The occurrence of "salmonella" in waste water from Danish slaughterhouses. A quantitative study.

L11 ANSWER 56 OF 85 MEDLINE  
T1 Alteration of low and high affinities of secreted and cell-bound antibodies during the anamnestic response of rabbits to "Salmonella" "senftenberg" microorganisms.

L11 ANSWER 57 OF 85 MEDLINE  
T1 Incidence of infections with "Salmonella" enteritidis serotypes in Black and Indian children. A 16-year survey.

L11 ANSWER 58 OF 85 MEDLINE  
T1 "Salmonella" serotypes encountered in animal feed additives in Lebanon.

L11 ANSWER 59 OF 85 MEDLINE  
T1 The effect of compounds which degrade hydrogen peroxide on the enumeration of heat-stressed cells of "Salmonella" "senftenberg" .

L11 ANSWER 60 OF 85 MEDLINE  
T1 [Isolation of R-phase strains of S. "senftenberg" from fish meal]. Proučavanje na izolirani od ribeno brašmo sfilamove S. "senftenberg" v R-faza.

L11 ANSWER 61 OF 85 MEDLINE  
T1 A survey of "normal" broiler mortality in East Anglia.

L11 ANSWER 62 OF 85 MEDLINE  
T1 Age-dependent resistance of chicken of "Salmonella" in vitro: antibacterial activity of lysed granule fraction of splenic adherent cells.

L11 ANSWER 63 OF 85 MEDLINE  
T1 Age-dependent resistance of chickens to "salmonella" in vitro: phagocytic and bactericidal activities of splenic phagocytes.

L11 ANSWER 64 OF 85 MEDLINE  
T1 Culture method for detection of "Salmonella" in dried active yeast: collaborative study.

L11 ANSWER 65 OF 85 MEDLINE  
T1 Minimal medium recovery of thermally injured "Salmonella" "senftenberg" 4969.

L11 ANSWER 66 OF 85 MEDLINE

T1 [Do immunization of rabbits by N-acetyl-galactosamine and a disaccharide linked to a protein produce anti-microbial antibodies (author's trans)]. Essai de production chez le lapin d'anticorps antimicrobiens par immunisation avec la N-acetyl-galactosamine et un dioside liés a une protaine.

L11 ANSWER 67 OF 85 MEDLINE  
T1 Destruction of "Salmonella" on poultry meat with lysozyme, EDTA, x-ray, microwave and chlorine.

L11 ANSWER 68 OF 85 MEDLINE  
T1 "Salmonella" survival on pecans as influenced by processing and storage conditions.

L11 ANSWER 69 OF 85 MEDLINE  
T1 Molecular immunological heterogeneity of the "Salmonella" zuendorf [1, 9, 12, (46), 27] cell-wall polysaccharides.

L11 ANSWER 70 OF 85 MEDLINE  
T1 [Changes in the specificity of antibodies appearing in the beginning of immunization by ""Salmonella" "senftenberg" (author's trans)]. Evolution de la specificite des anticorps apparaissant en debut d'immunisation par "Salmonella" "senftenberg" .

L11 ANSWER 71 OF 85 MEDLINE  
T1 Effect of water activity on heat survival of Staphylococcus aureus, "Salmonella" typhimurium and Salm. "senftenberg" .

L11 ANSWER 72 OF 85 MEDLINE  
T1 Inactivation of strains of "Salmonella" "senftenberg" by gamma irradiation.

L11 ANSWER 73 OF 85 MEDLINE  
T1 Viability of Staphylococcus aureus, "Salmonella" typhimurium and "Salmonella" "senftenberg" heated and recovered on a solid medium of controlled water activity.

L11 ANSWER 74 OF 85 MEDLINE  
T1 Epidemiological studies on "Salmonella" "senftenberg" . II. Infections in farm animals.

L11 ANSWER 75 OF 85 MEDLINE  
T1 Epidemiological studies on "Salmonella" "senftenberg" . I. Relations between animal foodstuff, animal and human isolations.

L11 ANSWER 76 OF 85 MEDLINE  
T1 "Salmonella" "senftenberg" in the Sunderland area

L11 ANSWER 77 OF 85 MEDLINE  
T1 Thermal inactivation of "Salmonella" "senftenberg" 775W in poultry meat.

L11 ANSWER 78 OF 85 MEDLINE  
T1 The effect of moisture and storage temperature on a "Salmonella" "senftenberg" 775W population in meat and bone meal

L11 ANSWER 79 OF 85 MEDLINE  
T1 Effect of pH and chelating agents on the heat resistance and viability of "Salmonella" typhimurium Tm-1 and "Salmonella" "senftenberg" 775W in egg white.

L11 ANSWER 80 OF 85 MEDLINE  
T1 Thermal resistance of "Salmonella" "senftenberg" 775W in dry animal feeds.

L11 ANSWER 81 OF 85 MEDLINE  
T1 Heat resistance of "Salmonella" : the uniqueness of "Salmonella" "senftenberg" 775W.

L11 ANSWER 82 OF 85 MEDLINE  
T1 Heat resistance of "Salmonella" typhimurium and "Salmonella" "senftenberg" 775W in milk chocolate.

L11 ANSWER 83 OF 85 MEDLINE  
T1 Thermal resistance of smooth and rough derivatives of "Salmonella" "senftenberg" 775 W.

L11 ANSWER 84 OF 85 MEDLINE  
T1 Initial evaluation of the effect of butylated hydroxytoluene upon "Salmonella" "senftenberg" 775W.

L11 ANSWER 85 OF 85 MEDLINE  
T1 Heat resistance of "Salmonella" typhimurium and "Salmonella" "senftenberg" 775 W in chicken meat.

L11 ANSWER 22 OF 85 MEDLINE AN 93210422 MEDLINE DN 93210422  
T1 "Salmonella"-induced enteritis: Clinical, serotypes and treatment.  
AU Ramadan F; Uhmri A G; Hablas R; Rizk M S  
CS Medical Department, Royal Commission Hospital, Yanbu, Kingdom of Saudi Arabia. .

DT Journal Article; (JOURNAL ARTICLE) LA English EM 199307

AB "Salmonella"-induced enteritis is a widespread cause of morbidity and mortality especially in developing countries. The frequency of different "Salmonella" serotypes in different areas varies according to time and locality. The prevalence of different "Salmonella" serotypes in Yanbu area was studied in 136 stool cultures from patients admitted with gastroenteritis, to the medical ward of Royal Commission Hospital in the period 1/6/1991 to 30/10/1991. Fifteen different "Salmonella" serotypes were determined among 31 positive "Salmonella" isolates and all were of the gastroenteric group, diarrhoeagenic but noninvasive. The most common serotype was S. typhimurium (45.16%) followed by S. enteritidis (9.62%) then S. virchow (6.46%). Other forms of "Salmonella" were isolated from one patient each: S. paratyphi B java, S. heidelberg, S. livingstone, S. infantis, S. bovis morbificans, S. corvallis, S. eastbourne, S. give, S. "senftenberg", S. poona, S. adelaide, and S. johannesburg. Saudi patients comprised about 71% and 29% were patients of four different nationalities. Antibiograms of these cultures proved to be all sensitive to norfloxacin with different forms of resistance to chloramphenicol, ampicillin and trimethoprim. Norfloxacin proved to be effective in the treatment of resistant forms of "Salmonella" with negligible side effects and wide safety range.

L13 ANSWER 1 OF 1 MEDLINE

T1 Development of a conjoint phage typing & biotyping schema for "Salmonella" "enterica" serovar "Senftenberg" (S. "senftenberg") & the correlation of biotypes with phage types.



FILE 'REGISTRY' ENTERED AT 12:11:54 ON 09 DEC 1997  
L1 477 S HYDRATASE?

FILE 'MEDLINE' ENTERED AT 12:12:06 ON 09 DEC 1997  
L2 1 S L1

FILE 'REGISTRY' ENTERED AT 12:12:32 ON 09 DEC 1997  
L3 581 S DEHYDRATASE?

FILE 'MEDLINE' ENTERED AT 12:12:50 ON 09 DEC 1997  
L4 6 S L3

FILE 'REGISTRY' ENTERED AT 12:13:58 ON 09 DEC 1997  
L5 1 S GLYCEROL DEHYDRATASE  
L6 2 S DIOL DEHYDRATASE

FILE 'MEDLINE' ENTERED AT 12:14:36 ON 09 DEC 1997  
L7 0 S L5  
L8 0 S L6  
E DEHYDRATASES/CT  
E E4  
E DEHYDRATASE/CT  
E DEHYDRATASE/CN  
E HYDRO LYASES  
E HYDRO LYASES/CT  
L9 2938 S E9  
L10 77716 S CLONING, MOLECULAR/CT  
L11 125 S L9 AND L10  
L12 37111 S KLEBSIELLA OR LACTOBACILLUS OR ENTEROBACTER OR CITROBACTER OR  
PELOBACTER OR ILIYOBACTER OR CLOSTRIDIUM  
L13 4 S L11 AND L12  
E GLYCEROL DEHYDRATASE  
E GLYCEROL DEHYDRATASE/CT  
E DIOL DEHYDRATASE  
E DIOL DEHYDRATASE/CT  
E DIOL DEHYDRATASE/CN  
E GLYCEROL DEHYDRATASE/CN  
L14 12 S E3  
L15 0 S L14 NOT L9  
L16 165 S L12 AND L9 NOT L13  
E KLEBSIELLA/CN  
E KLEBSIELLA/CT  
E L9  
E HYDRO LYASES/CT  
L17 291 S E22  
L18 7 S L17 AND L12  
L19 3 S L18 NOT L13

FILE 'SCISEARCH' ENTERED AT 12:32:16 ON 09 DEC 1997  
E SPRENGER G, 1989/RE  
E SPRENGER G A, 1989/RE  
L20 9 S E4

L4 ANSWER 1 OF 5 MEDLINE  
T1 Site-directed mutagenesis of monofunctional chorismate mutase engineered from the E. coli P-protein.

L4 ANSWER 2 OF 5 MEDLINE  
T1 Genetic aspects of aromatic amino acid biosynthesis in *Lactococcus lactis*.

L4 ANSWER 3 OF 5 MEDLINE  
T1 The *pheA*/*tyrA*/*aroF* region from *Erwinia herbicola*: an emerging comparative basis for analysis of gene organization and regulation in enteric bacteria.

L4 ANSWER 4 OF 5 MEDLINE  
T1 Loss of allelic control but retention of the bifunctional catalytic competence of a fusion protein formed by excision of 260 base pairs from the 3' terminus of *pheA* from *Erwinia herbicola*.

L4 ANSWER 5 OF 5 MEDLINE  
T1 Cloning, sequencing, and expression of the P-protein gene (*pheA*) of *Pseudomonas stutzeri* in *Escherichia coli*: implications for evolutionary relationships in phenylalanine biosynthesis.

L13 ANSWER 1 OF 4 MEDLINE AN 96422012 MEDLINE  
T1 Cloning, sequencing, and overexpression of the genes encoding coenzyme B12-dependent glycerol dehydratase of *Citrobacter freundii*.  
AU Seyfried M; Daniel R; Gottschalk G  
CS Institut für Mikrobiologie der Georg-August-Universität, Göttingen, Germany  
SO JOURNAL OF BACTERIOLOGY, (1996 Oct) 178 (19) 5793-6. Journal code: HH3. ISSN: 0021-9193. CY United States  
DT Journal Article; (JOURNAL ARTICLE) LA English FS Priority Journals OS GENBANK-U09771 EM 9701 EW 19970104  
AB The genes encoding coenzyme B12-dependent glycerol dehydratase of *Citrobacter freundii* were cloned and overexpressed in *Escherichia coli*. The B12-free enzyme was purified to homogeneity. It consists of three types of subunits whose N-terminal sequences are in accordance with those deduced from the open reading frames *dhaB*, *dhaC*, and *dhaE*, coding for subunits of 60 433 (alpha), 21 487 (beta), and 16 121 (gamma) Da, respectively. The enzyme complex has the composition  $\alpha_2\beta_2\gamma_2$ . Amino acid alignments with the subunits of the recently sequenced diol dehydratase of *Klebsiella oxytoca* (T. Tobimatsu, T. Hara, M. Sakaguchi, Y. Kishimoto, T. Wada, M. Isoda, T. Sakai, and T. Toraya, J. Biol. Chem. 270:7142-7148, 1995) revealed identities between 51.8 and 70.9%.  
CT Check Tags: Comparative Study Bacterial Proteins: BI, biosynthesis Bacterial Proteins: GE, genetics Bacterial Proteins: IP, isolation & purification Chromatography, Affinity Citrobacter freundii: EN, enzymology Citrobacter freundii: GE, genetics Citrobacter freundii: IP, isolation & purification Cobamides: ME, metabolism Escherichia coli: GE, genetics Escherichia coli: IP, isolation & purification Genes, Bacterial Hydro-Lyases: BI, biosynthesis Hydro-Lyases: GE, genetics Hydro-Lyases: IP, isolation & purification Molecular Sequence Data Protein Conformation Recombinant Proteins: BI, biosynthesis Recombinant Proteins: IP, isolation & purification Sequence Analysis, DNA Sequence Homology, Amino Acid Species Specificity  
CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.30 (glycerol dehydratase); 0 (Bacterial Proteins); 0 (Cobamides); 0 (Recombinant Proteins)

L13 ANSWER 2 OF 4 MEDLINE AN 96394290 MEDLINE  
T1 Cloning, sequencing, and high level expression of the genes encoding adenosylcobalamin-dependent glycerol dehydratase of *Klebsiella pneumoniae*.  
AU Tobimatsu T; Azuma M; Matsubara H; Takatori H; Niida T; Nishimoto K; Satoh H; Hayashi R; Toraya T  
CS Department of Bioscience and Biotechnology, Faculty of Engineering, Okayama University, Tushima-Naka, Okayama 700, Japan.  
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1996 Sep 13) 271 (37) 22352-7. Journal code: HIV. ISSN: 0021-9258. CY United States DT Journal Article; (JOURNAL ARTICLE) LA English FS Priority Journals Cancer Journals EM 9701 EW 19970104  
AB The *gid* genes encoding adenosylcobalamin-dependent glycerol dehydratase of *Klebsiella pneumoniae* were cloned by cross-hybridization with a DNA fragment of *Klebsiella oxytoca* diol dehydratase genes. Since the *Escherichia coli* clones isolated did not show appreciable enzyme activity, plasmids for high level expression of cloned genes were constructed. The enzyme expressed in *E. coli* was indistinguishable from the wild-type glycerol dehydratase of *K. pneumoniae* by the criteria of polyacrylamide gel electrophoresis, immunochemical, and catalytic properties. It was also shown that the recombinant functional enzyme consists of Mr 61 000, 22 000, and 16 000 subunits. Sequence analysis of the genes revealed four open reading frames separated by 2-12 bases. The sequential three open reading frames from the first to the third (*gidA*, *gidB*, and *gidC* genes) encoded polypeptides of 555, 194, and 141 amino acid residues with predicted molecular weights of 60 659 (alpha), 21 355 (beta), and 16 104 (gamma), respectively. High level expression of these three genes in *E. coli* produced more than 14-fold higher level of fully active apoenzyme than that in *K. pneumoniae*. It was thus concluded that these are the genes encoding the subunits of glycerol dehydratase. The deduced amino acid sequences of the three subunits were 71, 58, and 54% identical with those of the alpha, beta, and gamma subunits of diol dehydratase, respectively, but failed to show any apparent homology with other proteins.  
CT Check Tags: Support, Non-U.S. Gov't Amino Acid Sequence Base Sequence Cloning, Molecular DNA, Bacterial CH, chemistry Electrophoresis, Gel, Two-Dimensional Escherichia coli Gene Expression Regulation, Enzymologic Enzymology Hydro-Lyases: GE, genetics Hydro-Lyases: ME, metabolism Kiebsiella pneumoniae: EN, enzymology Molecular Sequence Data Plasmids: ME, metabolism Propanediol Dehydratase: CH, chemistry Propanediol Dehydratase: GE, genetics Propanediol Dehydratase: ME, metabolism Restriction Mapping Sequence Homology, Amino Acid Sequence Homology, Nucleic Acid  
CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.28 (Propanediol Dehydratase); EC 4.2.1.30 (glycerol dehydratase); 0 (DNA, Bacterial); 0 (Plasmids)

L13 ANSWER 3 OF 4 MEDLINE AN 93122543 MEDLINE  
T1 Growth temperature-dependent activity of glycerol dehydratase in *Escherichia coli* expressing the *Citrobacter freundii dha* regulon.  
AU Daniel R; Gottschalk G  
CS Institute für Mikrobiologie, Georg-August-Universität, Göttingen, FRG..



- SO FEMS MICROBIOLOGY LETTERS, (1992 Dec 15) 79 (1-3) 281-5. Journal code: F.M.L. ISSN: 0378-1097. CY Netherlands DT Journal, Article; (JOURNAL ARTICLE)  
LA English FS Priority Journals EM 9304  
AB Using the cosmid pWE15, a genomic library of **\*\*\*Citrobacter\*\*\*** freundii DNA in Escherichia coli ECL707 was prepared and screened for glycerol utilization. Six out of approximately 3000 clones were positive. One clone, harboring the recombinant cosmid pRO1, expressed glycerol dehydrogenase in high activity when grown at 28 degrees C but not at 37 degrees C. The growth temperature had little effect on the activity of the other enzymes encoded by the dha regulon. When the glycerol-containing medium was supplemented with cornitols, the recombinant E. coli strain produced 1,3-propanediol in high amounts at 28 degrees C.  
CT Check Tags: Support, Non-U.S. Govt. **\*\*\*Citrobacter freundii: EN, enzymology\*\*\*** **\*\*\*Citrobacter freundii: GE, genetics\*\*\***  
Cloning, Molecular **\*\*\*Escherichia coli: EN, enzymology\*\*\*** **Escherichia coli: GE, genetics**  
Genes, Bacterial **Propanediol: GE, metabolism** **\*\*\*Hydro-Lyases: GE, genetics\*\*\***  
metabolism **\*\*\*Propanediol: GE, metabolism\*\*\*** **Temperature**  
CN 504-63-2 (1,3-propanediol); 56-81-5 (Glycerol)  
RN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.30 (glycerol dehydratase); 0 (Propanediols)  
GEN dha
- L13 ANSWER 4 OF 4 MEDLINE AN 92412068 MEDLINE  
T1 Cloning and properties of a cyanide hydratase gene from the phytopathogenic fungus Gloeocercospora sorghi.  
AU Wang P; VanEten H D  
CS Department of Plant Pathology, University of Arizona, Tucson 85721..  
SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1992 Sep 16) 187 (2) 1048-54. Journal code: 9Y8. ISSN: 0006-291X. CY United States DT Journal, Article; (JOURNAL ARTICLE) LA English FS Priority Journals; Cancer Journals  
OS GENBANK-M89044; GENBANK-S41678; GENBANK-S41680; GENBANK-S41731; GENBANK-D10916; GENBANK-D10917; GENBANK-D10918; GENBANK-D10919; GENBANK-D10920 EM 9212  
AB The Cht gene encoding cyanide hydratase (CHT; EC 4.2.1.66), which detoxifies HCN and is thought to be important in fungal infection of cyanogenic plants, has been cloned from the phytopathogenic fungus Gloeocercospora sorghi. The gene was isolated by screening an expression library of G. sorghi using a CHT-specific antibody and using one of the positive cDNA clones as a probe in Southern hybridization to identify a 3.1 kb PstI genomic fragment. This PstI fragment expressed CHT activity when transformed into Aspergillus nidulans, a fungus that normally lacks CHT activity. Sequence analysis identified a single open reading frame of 1,107 base pairs which encodes a polypeptide of 40,904 daltons. The deduced amino acid sequence of CHT shares 36.5% identity to a nitrilase from the bacterium **\*\*\*Klebsiella\*\*\* pneumoniae subsp. ozaenae**.  
CT Check Tags: Comparative Study **Amino Acid Sequence** **Aminohydrolyases: CH, chemistry** **Aspergillus nidulans: GE, genetics**  
Base Sequence **Blotting, Southern** **DNA, CH, chemistry** **DNA: IP, isolation & purification** **DNA**  
Probes **\*\*\*Hydro-Lyases: CH, chemistry\*\*\*** **\*\*\*Hydro-Lyases: GE, genetics\*\*\*** **Hyphomycetes: EN, enzymology**  
Hyphomycetes: GE, genetics **\*\*\*Klebsiella pneumoniae: EN, enzymology\*\*\*** **Molecular Sequence Data** **Nucleic Acid Hybridization**  
Poly A; GE, genetics **Potassium Cyanide: PO, pharmacology** **RNA: GE, genetics** **Sequence Homology, Nucleic Acid** **Transcription, Genetic** **Transformation, Genetic** **Translation, Genetic**  
RN 151-50-8 (Potassium Cyanide); 24937-83-5 (Poly A); 63231-63-0 (RNA); 9007-49-2 (DNA)  
CN EC 3.5.4. (Aminohydrolyases); EC 3.5.5.1 (nitrilase); EC 4.2.1. (Hydro-Lyases); EC 4.2.1.66 (cyanide hydratase); 0 (DNA Probes); 0 (RNA, Messenger)
- L19 ANSWER 1 OF 3 MEDLINE  
T1 Analysis of acyl coenzyme A binding to the transcription factor FasR and identification of amino acid residues in the carboxyl terminus required for ligand binding.
- L19 ANSWER 2 OF 3 MEDLINE  
T1 The nucleotide sequence of genes involved in the leucine biosynthetic pathway of **\*\*\*Clostridium\*\*\* pasteurianum**.
- L19 ANSWER 3 OF 3 MEDLINE AN 90155202 MEDLINE  
T1 Anaerobic growth of Escherichia coli on glycerol by importing genes of the dha regulon from **\*\*\*Klebsiella\*\*\* pneumoniae**.  
AU Sprenger G A; Hammer B A; Johnson E A; Lin E C  
CS Department of Microbiology and Molecular Genetics, Harvard Medical School, Boston, MA 02115.  
NC 5-R01-GM11983 (NIGMS)  
SO JOURNAL OF GENERAL MICROBIOLOGY, (1989 May) 135 ( Pt 5) 1255-62. Journal code: 187. ISSN: 0022-1287. CY ENGLAND: United Kingdom  
DT Journal, Article; (JOURNAL ARTICLE) LA English FS Priority Journals EM 9005  
AB The dha regulon of **\*\*\*Klebsiella\*\*\* pneumoniae** specifying fermentative dissimilation of glycerol was mobilized by the broad-host-range plasmid RP4::mini Mu and introduced conjugatively into Escherichia coli. The recipient E. coli was enabled to grow anaerobically on glycerol without added hydrogen acceptors, although its cell yield was less than that of K. pneumoniae. The reduced cell yield was probably due to the lack of the coenzyme-B12-dependent glycerol dehydratase of the dha system. This enzyme initiates the first step in an auxiliary pathway for disposal of the extra reducing equivalents from glycerol. The lack of this enzyme would also account for the absence of 1,3-propanediol (a hallmark fermentation product of glycerol) in the spent culture medium. In a control experiment, a large quantity of this compound was detected in a similar culture medium following the
- growth of K. pneumoniae. The other three known enzymes of the dha system, glycerol dehydrogenase, dihydroxyacetone kinase and 1,3-propanediol oxidoreductase, however, were synthesized at levels comparable to those found in K. pneumoniae. Regulation of the dha system in E. coli appeared to follow the same pattern as in K. pneumoniae: the three acquired enzymes were induced by glycerol, catabolite repressed by glucose, and glycerol dehydrogenase was post-translationally inactivated during the shift from anaerobic to aerobic growth. The means by which the E. coli recipient can achieve redox balance without formation of 1,3-propanediol during anaerobic growth on glycerol remains to be discovered.  
CT Check Tags: Support, Non-U.S. Govt. Support, U.S. Govt. P.H.S. **Alcohol Oxidoreductases: GE, genetics** **Anaerobiosis** **Bacterial**  
Proteins: GE, genetics **Conjugation, Genetic** **Energy Metabolism** **Escherichia coli: GD, growth & development** **Escherichia coli: GE, genetics** **Escherichia coli: ME, metabolism** **Fermentation** **\*Genes, Bacterial** **Glycerin: ME, metabolism** **\*\*\*Hydro-Lyases: GE, genetics\*\*\*** **Hydro-Lyases: PH, physiology** **\*\*\*Klebsiella pneumoniae: GE, genetics\*\*\*** **Oxidation-Reduction**  
Phosphotransferases: GE, genetics **Plasmids** **Sugar Alcohol Dehydrogenases: GE, genetics**  
RN 56-81-5 (Glycerin)  
CN EC 1.1. (Alcohol Oxidoreductases); EC 1.1. (Sugar Alcohol Dehydrogenases); EC 1.1.1.6 (glycerol dehydrogenase); EC 1.1.1.77 (acetaldehyde reductase); EC 2.7 (Phosphotransferases); EC 2.7.1.29 (glycerone kinase); EC 4.2.1. (Hydro-Lyases); EC 4.2.1.30 (glycerol dehydratase); 0 (Bacterial Proteins); 0 (Plasmids)
- L20 ANSWER 1 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 Glycerol conversion to 1,3-propanediol by Clostridium pasteurianum: cloning and expression of the gene encoding 1,3-propanediol dehydrogenase
- L20 ANSWER 2 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 Structure and gene-polypeptide relationships of the region encoding glycerol diffusion facilitator (gpf) and glycerol kinase (gpk) of Pseudomonas aeruginosa
- L20 ANSWER 3 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 BIOCHEMICAL AND MOLECULAR CHARACTERIZATION OF THE OXIDATIVE BRANCH OF GLYCEROL UTILIZATION BY CITROBACTER-FREUNDII
- L20 ANSWER 4 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 CLONING AND NUCLEOTIDE-SEQUENCE OF THE GLD P GENE ENCODING SN-GLYCEROL-3-PHOSPHATE DEHYDROGENASE OF PSEUDOMONAS-AERUGINOSA
- L20 ANSWER 5 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 DIOL DEHYDRASE AND GLYCEROL DEHYDRASE, COENZYME B-12-DEPENDENT ISOZYMES
- L20 ANSWER 6 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 MAPPING AND CLONING OF GLDA, THE STRUCTURAL GENE OF THE ESCHERICHIA-COLI GLYCEROL DEHYDROGENASE
- L20 ANSWER 7 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 ANALYSIS OF THE ESCHERICHIA-COLI GENOME 4. DNA-SEQUENCE OF THE REGION FROM 89.2 TO 92.8 MINUTES
- L20 ANSWER 8 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 GROWTH TEMPERATURE-DEPENDENT ACTIVITY OF GLYCEROL DEHYDRATASE IN ESCHERICHIA-COLI EXPRESSING THE CITROBACTER-FREUNDII DHA REGULON
- L20 ANSWER 9 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R)  
T1 1,3-PROPANEDIOL PRODUCTION BY ESCHERICHIA-COLI EXPRESSING GENES FROM THE KLEBSIELLA-PNEUMONIAE-DHA REGULON
- L20 ANSWER 1 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R) AN 97-718159 SCISEARCH  
GA The Genuine Article (R) Number: XX393  
T1 Glycerol conversion to 1,3-propanediol by Clostridium pasteurianum: cloning and expression of the gene encoding 1,3-propanediol dehydrogenase  
AU Luers F; Seyfried M; Daniel R; Gottschalk G (Reprint)  
CS UNIV GOTTINGEN, INST MIKROBIOL, GRIEBBACHSTR 8, D-37077 GOTTINGEN, GERMANY (Reprint); UNIV GOTTINGEN, INST MIKROBIOL, D-37077 GOTTINGEN, GERMANY  
SO FEMS MICROBIOLOGY LETTERS, (15 SEP 1997) Vol. 154, No. 2, pp. 337-345. Publisher: ELSEVIER SCIENCE BV, PO BOX 211, 1000 AE AMSTERDAM, NETHERLANDS.  
ISSN: 0378-1097. DT Article; Journal FS LIFE LA English REC Reference Count: 34  
AB When grown on glycerol as sole carbon and energy source, cell extracts of Clostridium pasteurianum exhibited activities of glycerol dehydrogenase, dihydroxyacetone kinase, glycerol dehydratase and 1,3-propanediol dehydrogenase. The genes encoding the latter two enzymes were cloned by colony hybridization using the dha1 gene of Citrobacter freundii as heterologous DNA probe and expressed in Escherichia coli. The native molecular mass of 1,3-propanediol dehydrogenase (DhaT) is 440 000 Da. The dha1 gene of C. pasteurianum was subcloned and its nucleotide sequence (1158 bp) was determined. The deduced gene product (41 776 Dal revealed high similarity to DhaT of C. freundii (80.5% identity; 89.8% similarity).  
CT MICROBIOLOGY  
ST Author Keywords: Clostridium pasteurianum; 1,3-propanediol dehydrogenase; 1,3-propanediol; glycerol fermentation; type III alcohol dehydrogenase; glycerol dehydratase

STP KeyWords Plus (R): ESCHERICHIA-COLI; ALCOHOL-DEHYDROGENASE; CITROBACTER-FREUNDII; MOLECULAR CHARACTERIZATION; KLEBSIELLA-PNEUMONIAE; ZYMONOMAS-MOBILIS; SEQUENCE-ANALYSIS; OHA REGULON; PROTEINS; OVEREXPRESSION; RF 95-0636 001; 11-BETA-HYDROXYSTEROID DEHYDROGENASE; FETAL ORIGINS OF CORONARY HEART-DISEASE; APPARENT MINERALOCORTICOID EXCESS SYNDROMES 95-3190 001; INCREASED ABUNDANCE OF SPECIFIC SKELETAL-MUSCLE PROTEIN-TYROSINE PHOSPHATASES; ALPHA-B-CRYSTALLIN EXPRESSION 95-3375 001; THERMUS STRAINS; DNA RELATEDNESS; GENUS AEROMONAS; EMENDED DESCRIPTION OF CAMPYLOBACTER-HYOMINTESINALIS; POLYPHASIC TAXONOMY 95-5061 001; STRUCTURAL GENE, GLYC-DEPENDENT REGULATION OF BACILLUS-SUBTILIS GLUTAMATE SYNTHASE EXPRESSION; ARABIDOPSIS TYPE-1 PROTEIN PHOSPHATASE

RE Referenced Author | Year | Vol | Pg | Referenced Work  
(RAU) [(RPV)](RVL)](RPG)] (RWK)

=====

ABBADANDAUSSIS 1996 1142 1149 JMICROBIOL-UK  
ANDERSSON L O 1972 20 199 JFEBS LETT  
AUSUBEL F M 1987 1 1 CURRENT PROTOCOLS MO  
BAROCHA 1991 119 1241 JNUCLEIC ACIDS RES  
BOENIGK R 1991 1 1 THESIS G AUGUST U GO  
BRADFORD M M 1976 172 1248 JAVAL BIOCHEM  
CONWAY T 1987 169 2591 J BACTERIOL  
CONWAY T 1989 1171 3754 J BACTERIOL  
DABROCK B 1992 158 1123 JAPPL ENVIRON MICROB  
DANIEL R 1992 100 281 JFEMS MICROBIOL LETT  
DANIEL R 1995 177 12151 J BACTERIOL  
DANIEL R 1995 1177 4392 J BACTERIOL  
DEVRIES G E 1992 1174 15346 J BACTERIOL  
FISCHER R J 1993 1175 16659 J BACTERIOL  
GOODLOVE P E 1989 185 1209 JGENE  
HEYDRICKX M 1991 134 1637 JAPPL MICROBIOL BIOT  
HOMANN T 1990 133 1121 JAPPL MICROBIOL BIOT  
JOHNSON E A 1984 1160 155 J BACTERIOL  
KELL D B 1981 199 181 BIOCHEM BIOPH RES CO  
KESSLER D 1992 267 118073 J BIOL CHEM  
NARMUR J 1981 13 1208 J MOL BIOL  
REID M F 1984 120 113 JCRIT REV MICROBIOL  
RUCH F E 1977 119 150 J BACTERIOL  
RUCH F E 1977 174 15463 J NATL ACAD SCI USA  
SEYFRIED M 1986 1178 15793 J BACTERIOL  
SOHLING B 1996 1178 1871 J BACTERIOL  
SPRENGER G A 1989 1135 11255 J GEN MICROBIOL  
STOILJKOVIC I 1995 1177 1367 J BACTERIOL  
TORAYA T 1977 1252 1963 J BIOL CHEM  
TSE P 1988 1110 11295 J AM CHEM SOC  
WALTER K A 1992 1174 17149 J BACTERIOL  
WIERENGA R K 1985 124 11346 J BIOCHEMISTRY-US  
WILLAMSON V M 1987 1209 1374 J MOL GEN GENET  
YOUNGLESON J S 1988 178 1355 JGENE

L20 ANSWER 3 OF 9 SCISEARCH COPYRIGHT: 1997 ISI (R) AN 95 524431 SCISEARCH  
GA The Genuine Article (R) Number: RL828  
TI BIOCHEMICAL AND MOLECULAR CHARACTERIZATION OF THE OXIDATIVE BRANCH OF GLYCEROL UTILIZATION  
BY CITROBACTER-FREUNDII  
AU DANIEL R; STUERTZ K; GOTTSCALK G (Reprint)  
CS UNIV GOTTINGEN, INST MIKROBIOL, GRIEBBACHSTR 8, D-37077 GOTTINGEN, GERMANY (Reprint); UNIV  
GOTTINGEN, INST MIKROBIOL, D-37077 GOTTINGEN, GERMANY CYA GERMANY  
SO JOURNAL OF BACTERIOLOGY, (AUG 1995) Vol. 177, No. 15, pp. 4392-4401. ISSN: 0021-9193. DT Article; Journal  
FS LIFE A ENGLISH REC Reference Count: 58  
AB Glycerol dehydrogenase (EC 1.1.1.6) and dihydroxyacetone kinase (EC 2.7.1.29) were purified from Citrobacter freundii.  
The dehydrogenase is a hexamer of a polypeptide of 43,000 Da. The enzyme exhibited a rather broad substrate specificity, but  
glycerol was the preferred substrate in the physiological direction. The apparent K(m)s of the enzyme for glycerol and NAD(+) were 1.27 mM and 57 mu M, respectively. The kinase is a dimer of a polypeptide of 57,000 Da. The enzyme was highly specific  
for the substrates dihydroxyacetone and ATP; the apparent K(m)s were 30 and 70 mu M, respectively. The DNA region which  
contained the genes encoding glycerol dehydrogenase (dhad) and dihydroxyacetone kinase (dhak) was cloned and sequenced.  
Both genes were identified by N-terminal sequence comparison. The deduced dhad gene product (365 amino acids) exhibited  
high degrees of homology to glycerol dehydrogenases from other organisms and less homology to type III alcohol  
dehydrogenases, whereas the dhak gene product (552 amino acids) revealed no significant homology to any other protein in the  
databases. A large gene (dhakR) of 1,929 bp was found downstream from dhad. The deduced gene product (641 amino acids)  
showed significant similarities to members of the sigma(54) bacterial enhancer-binding protein family.  
CC MICROBIOLOGY

STP KeyWords Plus (R): ACTIVATED ALCOHOL-DEHYDROGENASE; METAL DISSOCIATION-CONSTANTS; ESCHERICHIA-COLI;  
KLEBSIELLA-PNEUMONIAE; ZYMONOMAS-MOBILIS; SACCHAROMYCES-CEREVISIAE; NUCLEOTIDE-SEQUENCE; OHA REGULON;  
BACILLUS-STEARTHERMOPHILUS; REGULATORY GENE  
RF 93-4847 004; HETEROLOGOUS EXPRESSION; CHROMOSOMAL DNA: GENE ENCODING METHYLMALONYL-COENZYME-A MUTASE  
93-7500 002; PROTEIN PHOSPHATASE-1; PHOTOTROPHIC BACTERIUM RHODOBACTER-CAPSULATUS EIF1; CALF UTERUS 93-3088  
001; RAT MUSCLE; PROTEIN PHOSPHATASE-1; MAJOR GLUTATHIONE TRANSFERASE 93-6277 001; ESCHERICHIA-COLI  
MESSENGER-RNA PROMOTER SEQUENCES; TRANSCRIPTION INITIATION; EXPRESSION OF THE CELLULOONOMAS-FLAVIGENA CELL-  
ASSOCIATED AMYLASE GENE 93-7923 001; SULFATE-REDUCING BACTERIUM; ANAEROBIC DEGRADATION; METHANE FORMATION

RE Referenced Author | Year | Vol | Pg | Referenced Work  
(RAU) [(RPV)](RVL)](RPG)] (RWK)

=====

ARNOLD W 1988 1203 1715 J MOL BIOL  
BAROCH A 1991 119 12241 JNUCLEIC ACIDS RES  
BLATTNER F R 1993 121 15408 JNUCLEIC ACIDS RES  
BLUM H 1987 18 153 J ELECTROPHORESIS  
BOENIGK R 1993 138 1463 JAPPL MICROBIOL BIOT  
BRADFORD M M 1976 172 1248 JAVAL BIOCHEM  
CONWAY T 1987 169 2591 J BACTERIOL  
CONWAY T 1989 1171 3754 J BACTERIOL  
DANIEL R 1992 100 281 JFEMS MICROBIOL LETT  
DANIEL R 1995 177 12151 J BACTERIOL  
DEVEREUX J 1984 112 1387 JNUCLEIC ACIDS RES  
DEVRIES G E 1992 1174 15346 J BACTERIOL  
DREWKE C 1988 1950 154 J BIOCHIM BIOPHYS ACTA  
FISCHER R J 1993 1175 16659 J BACTERIOL  
FORAGER R G 1982 1151 1591 J BACTERIOL  
FRY D C 1986 183 1907 J NATL ACAD SCI USA  
GOODLOVE P E 1989 185 1209 JGENE  
HAWLEY D K 1993 111 12237 JNUCLEIC ACIDS RES  
HOMANN T 1990 133 1121 JAPPL MICROBIOL BIOT  
INOUE S 1988 166 1301 JGENE  
JOHNSON E A 1984 1160 155 J BACTERIOL  
KESSLER D 1992 267 118073 J BIOL CHEM  
KRUGER N 1992 1174 14391 J BACTERIOL  
KYHSENDERSEN J 1984 110 1203 J BIOCHEM BIOPH METH  
LAEMMLI U K 1970 1270 1680 JNATURE  
MALLINDER P R 1992 1110 19 JGENE  
MARCK C 1988 116 1829 JNUCLEIC ACIDS RES  
MARTIN R G 1981 1236 11372 J BIOL CHEM  
MORETTE E 1993 1175 16067 J BACTERIOL  
PETTIGREW D W 1988 1263 1135 J BIOL CHEM  
PFENNIG N 1986 155 1245 JARCH MIKROBIOL  
RAMAKRISHNAN G 1990 187 12369 J NATL ACAD SCI USA  
REID M F 1984 120 113 JCRIT REV MICROBIOL  
RUCH F E 1974 119 150 J BACTERIOL  
RUCH F E 1980 1141 1077 J BACTERIOL  
SAMBROOK J 1989 1 1 JIMOL CLONING LABORATO  
SANGER F 1977 174 15463 J NATL ACAD SCI USA  
SHINE J 1974 171 1342 J NATL ACAD SCI USA  
SHINGLER V 1993 1175 11586 J BACTERIOL  
SIEGEL L M 1966 1112 1346 J BIOCHIM BIOPHYS ACTA  
SOHLING B 1 1 JNPUB  
SPENCER P 1989 1994 1270 J BIOCHIM BIOPHYS ACTA  
SPRENGER G A 1989 1135 11255 J GEN MICROBIOL <--  
SRIDHARA S 1969 198 187 J BACTERIOL  
SUTCLIFFE J G 1979 143 177 JCOLD SPRING HARB SYM  
TANG C T 1979 1140 1182 J BACTERIOL  
THORNER J W 1991 1246 13885 J BIOL CHEM  
TONG I T 1991 167 13541 JAPPL ENVIRON MICROB  
TORAYA T 1977 1252 1963 J BIOL CHEM  
TSE P 1988 1110 11295 J AM CHEM SOC  
TSE P 1989 1111 18703 J AM CHEM SOC  
VIERA J 1982 119 1259 JGENE  
WALTER K A 1992 1174 17149 J BACTERIOL  
WANG A Y 1992 131 111020 BIOCHEMISTRY-US  
WILLAMSON V M 1987 1209 1374 J MOL GEN GENET  
YAMADA H 1982 146 12333 JAGR BIOL CHEM TOKYO  
YANISCH-PERRON C 1985 133 1103 JGENE  
YOUNGLESON J S 1988 178 1355 JGENE

GA The Genuine Article (R) Number: B291T  
 TI DIOL DEHYDRASE AND GLYCEROL DEHYDRASE, COENZYME B-12-DEPENDENT ISOZYMES  
 AU TORAYA T (Reprint)  
 CS OKAYAMA UNIV, FAC ENGN, DEPT BIOTECHNOL, 3-1-1 TSUSHIMA NAKA, OKAYAMA 700, JAPAN (Reprint)  
 CYA JAPAN  
 SO METAL IONS IN BIOLOGICAL SYSTEMS, (1994) Vol. 30, pp. 217-254. ISSN: 0161-5149. DT General Review; Journal  
 LA ENGLISH REC Reference Count: 110  
 CC CHEMISTRY, INORGANIC & NUCLEAR; BIOLOGY, MISCELLANEOUS; BIOCHEMISTRY & MOLECULAR BIOLOGY; BIOPHYSICS  
 STP KeyWords Plus (R): BOND-DISSOCIATION ENERGY; CARBON-COBALT BOND; CO-CARBOND; KLEBSIELLA-PNEUMONIAE; CHEMICAL  
 MODIFICATION; ESCHERICHIA-COLI; DHA REGULON; D-RIbose; ADENOSYL-COBALAMIN; ENZYME  
 RE

Referenced Author [Year] [VOL] [PG] [Referenced Work]  
 (RAU) [RPV] (RV) [RPG] (RWK)

=====

ABELES R H 1976 9 114 J ACCOUNTS CHEM RES  
 ABELES R H 1964 112 695 J ANN NY ACAD SCI  
 ABELES R H 1960 141 531 J BIOCHEM BIOPHYS ACTA  
 ABELES R H 1971 15 1481 J ENZYMES  
 ABELES R H 1961 126 12347 J BIOL CHEM  
 ABELES R H 1966 1241 1245 J BIOL CHEM  
 ABELES R H 1966 9 1686 J METHOD ENZYMOL  
 ABELES R H 1979 1 373 J ITAMIN B12  
 ANTON D L 1980 102 12715 J AM CHEM SOC  
 ANTON D L 1980 255 14507 J BIOL CHEM  
 BABIOR B M 1988 11 121 J BIOFACTORS  
 BABIOR B M 1974 1249 11689 J BIOL CHEM  
 BACHOVCHIN W W 1977 116 1082 J BIOCHEMISTRY-US  
 BACHOVCHIN W W 1978 117 12218 J BIOCHEMISTRY-US  
 BONT J A M 1982 714 1465 J BIOCHEM BIOPHYS ACTA  
 BROWNSTEIN A M 1961 126 11199 J BIOL CHEM  
 COCKLE S A 1972 194 1275 J AM CHEM SOC  
 DANIEL R 1992 1100 1281 J FEMS MICROBIOL LETT  
 EAGAR R G 1975 114 15523 J BIOCHEMISTRY-US  
 EBERHARD G 1988 1369 1091 J BIOL CHEM HOPPESEYLE  
 ESENBERG M K 1971 193 11242 J AM CHEM SOC  
 FINKE R G 1984 154 11 J COORDIN CHEM REV  
 FINLAY T H 1972 1247 14197 J BIOL CHEM  
 FINLAY T H 1973 1248 1285 J BIOL CHEM  
 FORAGE R G 1979 1569 1249 J BIOCHEM BIOPHYS ACTA  
 FORAGE R G 1982 1149 1413 J BACTERIOL  
 FORAGE R G 1982 1151 1591 J BACTERIOL  
 FREY P A 1967 129 1873 J BIOCHEM BIOPH RES CO  
 FREY P A 1970 192 14488 J AM CHEM SOC  
 FREY P A 1966 1241 12732 J BIOL CHEM  
 FREY P A 1967 1242 15369 J BIOL CHEM  
 GOLDING B T 1982 11 1543 J B12  
 HALPERN J 1984 106 18317 J AM CHEM SOC  
 HARTMANIS M G N 1986 1245 1144 J ARCH BIOCHEM BIOPHYS  
 HAY B P 1986 108 14820 J AM CHEM SOC  
 HAY B P 1987 109 18012 J AM CHEM SOC  
 HONDA S 1980 1143 11458 J BACTERIOL  
 HOSOI N 1978 156 1566 J FERMENT TECHNOL  
 ICHIKAWA M 1988 1952 1191 J BIOCHEM BIOPHYS ACTA  
 ISHIDA A 1993 132 11535 J BIOCHEMISTRY-US  
 ISHIDA A 1993 139 1115 J NUTR SCI VITAMINOL  
 JENSEN F R 1975 142 1315 J METHOD ENZYMOL  
 JOHNSON B C 1975 142 1315 J METHOD ENZYMOL  
 KROUWER J S 1980 1612 1153 J BIOCHEM BIOPHYS ACTA  
 KUNO S 1980 1205 1240 J ARCH BIOCHEM BIOPHYS  
 KUNO S 1981 1210 1474 J ARCH BIOCHEM BIOPHYS  
 KUNO S 1981 1211 1722 J ARCH BIOCHEM BIOPHYS  
 KUNO S 1980 1277 1211 J ARCH BIOCHEM BIOPHYS  
 LEE H A 1963 1238 12367 J BIOL CHEM  
 MCGEE D E 1982 1108 1547 J BIOCHEM BIOPH RES CO  
 MCGEE D E 1981 120 14293 J BIOCHEMISTRY-US  
 MOORE K W 1979 187 11052 J BIOCHEM BIOPH RES CO  
 OBRADORS N 1988 1170 12159 J BACTERIOL  
 OCHIAI E I 1975 137 1351 J INORG NUCL CHEM  
 PAWELKIEWICZ J 1965 112 1207 J ACTA BIOCHEM POL  
 PAWELKIEWICZ J 1964 1112 1703 J ANN NY ACAD SCI

L20 ANSWER 8 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R) AN 93-17810 SCISEARCH  
 GA The Genuine Article (R) Number: KE540  
 TI GROWTH TEMPERATURE-DEPENDENT ACTIVITY OF GLYCEROL DEHYDRATASE IN ESCHERICHIA-COLI  
 EXPRESSING THE CITROBACTER-FREUNDII DHA REGULON  
 AU DANIEL R; GOTTSCHALK G (Reprint)  
 CS UNIV GOTTINGEN INST MIKROBIOL, GRIEBACHSTR 8, W-3400 GOTTINGEN, GERMANY CYA GERMANY  
 SO FEMS MICROBIOLOGY LETTERS, (15 DEC 1992) Vol. 100, No. 1-3, pp. 281-285. ISSN: 0378-1097. DT Article;  
 Journal FS LIFE LA ENGLISH REC Reference Count: 13  
 AB Using the cosmid pWE15, a genomic library of Citrobacter freundii DNA in Escherichia coli ECL707 was prepared and screened for glycerol utilization. Six out of approximately 3000 clones were positive. One clone, harboring the recombinant cosmid pRD1, expressed glycerol dehydratase in high activity when grown at 28-degrees-C but not at 37-degrees-C. The growth temperature had little effect on the activity of the other enzymes encoded by the dha regulon. When the glycerol-containing medium was supplemented with cornoids, the recombinant E. coli strain produced 1,3-propanediol in high amounts at 28-degrees-C.

CC MICROBIOLOGY

ST Author keywords: CITROBACTER-FREUNDII; ESCHERICHIA-COLI EQ.707; GLYCEROL DEHYDRATASE; 1,3-PROPANEDIOL; GLYCEROL FERMENTATION; DHA REGULON  
STP KeyWords Plus (R): KLEBSIELLA-PNEUMONIAE; ANAEROBIC GROWTH; GENES RF 92.3056 001; UPTAKE OF SURFACTANT PROTEIN-B; CASEIN KINASE-II; CATALYTIC SUBUNITS 92.4812 001; PUTATIVE ANAEROBIC COPROPORPHYRINOGEN-II OXIDASE IN RHODOBACTER-SPHAEROIDES; TRANSCRIPTIONAL REGULATORY ELEMENT; FUNCTIONAL EXPRESSION RE Referenced Author

[Year] [VOL] [PG] [Referenced Work] (RAU) [RPY] [RVL] [RPG] (RWK)

=====

BOENICK R	1991	1	THESIS GEORG AUGUST
BRADFORD M M	1976	172	248 JANAL BIOCHEM
EGGSTEN M	1974	1	METHODEN ENZYMATISCH
JETER R M	1984	159	206 J BACTERIOL
JOHNSON E A	1984	1160	155 J BACTERIOL
MARMUR J	1961	13	208 J MOL BIOL
PFENNIG N	1966	155	1245 JARCH MIKROBIOL
RUCH F E	1974	1119	150 J BACTERIOL
SAMBRON J	1989	1	J MOL CLONING
SPRENGER G A	1989	1135	1255 J GEN MICROBIOL
TONG I T	1991	157	3541 APPL ENVIRON MICROB
TORAYA T	1977	176	1285 JEUR J BIOCHEM
TORAYA T	1977	252	963 J BIOL CHEM

L20 ANSWER 9 OF 9 SCISEARCH COPYRIGHT 1997 ISI (R) AN 91:670800 SCISEARCH

GA The Genuine Article (R) Number: GT942  
TI 1,3-PROPANEDIOL PRODUCTION BY ESCHERICHIA-COLI EXPRESSING GENES FROM THE KLEBSIELLA-PNEUMONIAE-DHA REGULON  
AU TONG I T; LIAO H H; CAMERON D C (Reprint)  
CS UNIV WISCONSIN DEPT CHEM ENGN 1415 JOHNSON DR, MADISON, WI, 53706;  
UNIV WISCONSIN, CTR BIOTECHNOL, MADISON, WI, 53705 CYA USA  
SO APPLIED AND ENVIRONMENTAL MICROBIOLOGY, (1991) Vol. 57, No. 12, pp. 3541-3546. DT Article; Journal FS  
LIFE, AGRILA ENGLISH REC Reference Count: 33

AB The dha regulon in Klebsiella pneumoniae enables the organism to grow anaerobically on glycerol and produce 1,3-propanediol (1,3-PD). Escherichia coli, which does not have a dha system, is unable to grow anaerobically on glycerol without an exogenous electron acceptor and does not produce 1,3-PD. A genomic library of K. pneumoniae ATCC 25955 constructed in E. coli AG1 was enriched for the ability to grow anaerobically on glycerol and dihydroxyacetone and was screened for the production of 1,3-PD. The cosmid pTC1 (42.5 kb total with an 18.2-kb major insert) was isolated from a 1,3-PD-producing strain of E. coli and found to possess enzymatic activities associated with four genes of the dha regulon: glycerol dehydratase (dhaB), 1,3-PD oxidoreductase (dhaT), glycerol dehydrogenase (dhaD), and dihydroxyacetone kinase (dhaK). All four activities were inducible by the presence of glycerol. When E. coli AG1/pTC1 was grown on complex medium plus glycerol, the yield of 1,3-PD from glycerol was 0.46 mol/mol. The major fermentation by-products were formate, acetate, and D-lactate. 1,3-PD is an intermediate in organic synthesis and polymer production. The 1,3-PD fermentation provides a useful model system for studying the interaction of a biochemical pathway in a foreign host and for developing strategies for metabolic pathway engineering.  
CC MICROBIOLOGY; BIOTECHNOLOGY & APPLIED MICROBIOLOGY STP KeyWords Plus (R): GLYCEROL; DISSIMILATION DEHYDRATASES; COENZYME; KINASE RF 91:515 001; PHYSICAL MAP OF THE ESCHERICHIA-COLI CHROMOSOME; METZ GENE ENCODING TRANSFER-RNA MET F1; ASC (FORMERLY SAC) OPERON RE Referenced Author [Year] [VOL] [PG] [Referenced Work] (RAU) [RPY] [RVL] [RPG] (RWK)

=====

AUSUBEL F M	1987	1	CURRENT PROTOCOLS MO
BAILEY J E	1991	252	11668 SCIENCE
CAMERON D C	1990	1	J P CORN UT C ST LOU
COZZARELLIN R	1965	190	11325 J BACTERIOL
DANIELS L	1975	190	11325 APPL BACTERIOL
ELMR	1980	119	1425 JULLMANNS ENCY TECHNI
FORAGER G	1979	563	1249 BIOCHIM BIOPHYS ACTA
FORAGER G	1982	149	413 J BACTERIOL
FORAGER G	1982	1151	1591 J BACTERIOL
HONDA S	1980	1143	11458 J BACTERIOL
JOHNSON E A	1984	1160	155 J BACTERIOL
JOHNSON E A	1985	1164	1479 J BACTERIOL
JOHNSON E A	1987	1169	12050 J BACTERIOL
LIN E C C	1976	130	1535 JANNU REV MICROBIOL
LIN E C C	1980	1235	11820 J BIOL CHEM
LJUNGDAHL L G	1986	1	JMANUAL IND MICROBIOL
MACQUITT J J	1988	132	111 JACS SYM SER
MORRIS D W	1981	11	127 IDNA CELL BIOL
PAWELKIEWICZ J	1985	112	1207 JACTA BIOCHIM POL
RICHEY D P	1972	1112	1784 J BACTERIOL

RUCH F E 1974 1119 150 J BACTERIOL  
SCHNEIDER Z 1966 113 311 JACTA BIOCHIM POL  
SPRENGER G A 1989 135 1255 J GEN MICROBIOL  
SRIDHARA S 1969 198 187 J BACTERIOL  
STROINSKI A 1974 1162 321 JARCH BIOCHEM BIOPHYS  
TEMPEST D W 1987 1 1797 ESCHERICHIA COLI SAL  
TORAYA T 1982 1 233 112 BIOCH MED  
TORAYA T 1977 1252 963 J BIOL CHEM  
ZWAIG N 1970 1102 1753 J BACTERIOL

STN Patent No. [Year] [Ref. Inventor/Assignee] [Type] [Ref. Patent No. (RPN)] [RPY] (RIN) I I (RPN)

US 2465319 1949 WHINFIELD J R I JUS 2465319  
US 4873379 1989 MURPHY M A I JUS 4873379  
US 4937314 1990 GREENE R N I JUS 4937314

DIALOG INFORMATION SERVICES

09dec97 12:47:19 User208600 Session D1120.1

File 301:CHEMNAME(R) 1957-1997/Nov (c) 1997 Amer.Chem.Soc.

S1 1 GLYCEROL(W)DEHYDRATASE  
S2 1 DIOL(W)DEHYDRATASE

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1966-1997/Dec W4 (c) format only 1997 Knight-Ridder Info  
File 5:BIOSIS PREVIEW(S)(R) 1969-1997/Dec W1 (c) 1997 BIOSIS  
File 73:EMBASE 1974-1997/Nov W3 (c) 1997 Elsevier Science B.V.  
File 351:DERWENT WPI 1963-1997/UD=9748;UP=9745;UM=9743 (c)1997 Derwent Info Ltd

Set Items Description

S1 240 ADENOSYLCOBALAMIN(DEPENDENT(DIOL(D)DEHYDRASE + COENZYME(I) B12(D)DEPENDENT(DIOL(D)DEHYDRASE + COENZYME(B12(D)DEPENDENT(DIOL(D)DEHYDRATASE + DEHYDRATASE(DIOL + DIOL(D)DEHYDRASE + DIOL(D)DEHYDRATASE + MESO(12(3)BUTANEDIOL(D)DEHYDRASE  
S2 117 PROPANEDIOL(D)DEHYDRASE + PROPANEDIOL(D)DEHYDRATASE + 1(2)PROPANEDIOL(D)DEHYDRATASE  
S3 280 S1:S2  
S4 191 COENZYME(B12(D)DEPENDENT(DIOL(D)DEHYDRASE + COENZYME(I) GLYCEROL(D)DEHYDRASE + GLYCEROL(D)DEHYDRATASE  
S5 156571 KLEBSIELLA OR CITROBACTER OR LACTOBACILLUS OR ENTEROBACTEROR PELOBACTER OR ILYOBIOTER OR CLOSTRIDIUM  
S6 138 S3 AND S5  
S7 105 RD (unique items)  
S8 90 S4 AND S5 NOT S6  
S9 64 RD (unique items)

7/6/1 (Item 1 from file: 155) 091142159 97296406  
Kinetic investigations with inhibitors that mimic the posthomolysis intermediate in the reactions of coenzyme-B12-dependent glycerol dehydratase and diol dehydratase.

7/6/2 (Item 2 from file: 155) 08960494 97157051  
An electron paramagnetic resonance study on the mechanism-based inactivation of adenosylcobalamin-dependent diol dehydratase by glycerol and other substrates.

7/6/3 (Item 3 from file: 155) 08791004 96364290  
Cloning, sequencing, and high level expression of the genes encoding adenosylcobalamin-dependent glycerol dehydratase of Klebsiella pneumoniae.

7/6/4 (Item 4 from file: 155) 08790962 96422012  
Cloning, sequencing, and overexpression of the genes encoding coenzyme B12-dependent glycerol dehydratase of Citrobacter freundii.

- 7/6/5 (Item 5 from file: 155) 08213743 95271362  
Molecular cloning, sequencing, and expression of the genes encoding adenosylcobalamin-dependent diol dehydratase of *Klebsiella oxytoca*.
- 7/6/6 (Item 6 from file: 155) 07662694 94015511  
Importance of the nucleotide loop moiety coordinated to the cobalt atom of adenosylcobalamin for coenzymic function in the diol dehydratase reaction.
- 7/6/7 (Item 7 from file: 155) 07487018 93160191  
Adenosylcobinamide methyl phosphate as a pseudocoenzyme for diol dehydratase.
- 7/6/8 (Item 8 from file: 155) 06454075 90165470  
Essential histidine residues in coenzyme B12-dependent diol dehydratase: dye-sensitized photooxidation and ethoxycarbonylation.
- 7/6/9 (Item 9 from file: 155) 06216664 87092400  
Solubilization of a membrane-bound diol dehydratase with retention of EPR  $g = 2.02$  signal by using 2-(*N*-cyclohexylamino)ethanesulfonic acid buffer.
- 7/6/10 (Item 10 from file: 155) 06148838 87265998  
Re-investigation of the protein structure of coenzyme B12-dependent diol dehydratase.
- 7/6/11 (Item 11 from file: 155) 06130099 86129441  
Diol metabolism and diol dehydratase in *Clostridium glycolicum*.
- 7/6/12 (Item 12 from file: 155) 06086412 88107822  
Roles of the beta-D-ribofuranose ring and the functional groups of the D-ribose moiety of adenosylcobalamin in the diol dehydratase reaction.
- 7/6/13 (Item 13 from file: 155) 05575726 86207091  
[Studies on the biological function of the nucleotide base of vitamin B12] Untersuchungen zur biologischen Funktion der Nucleotidbase von Vitamin B12.
- 7/6/14 (Item 14 from file: 155) 05554022 88198006  
Anaerobic metabolism of the L-ribose fermentation product 1,2-propanediol in *Salmonella typhimurium*.
- 7/6/15 (Item 15 from file: 155) 05314974 87250467  
Activation and cleavage of the carbon-cobalt bond of adenosylcobalamin by diol dehydratase.
- 7/6/16 (Item 16 from file: 155) 05279470 86250875  
The synthesis of adenosine-modified analogs of adenosylcobalamin and their coenzymic function in the reaction catalyzed by diol dehydratase.
- 7/6/17 (Item 17 from file: 155) 04855866 86049396  
The binding site for the adenosyl group of coenzyme B12 in diol dehydratase.
- 7/6/18 (Item 18 from file: 155) 04410929 80182104  
The synthesis and properties of four spin-labeled analogs of adenosylcobalamin.
- 7/6/19 (Item 19 from file: 155) 03879299 82066866  
Chemical modification of coenzyme B12-dependent diol dehydratase with pyridoxal 5-phosphate: lysyl residue essential for interaction between two components of the enzyme.
- 7/6/20 (Item 20 from file: 155) 03841000 83074700  
Diol dehydratase: N-terminal amino acid sequences and subunit stoichiometry.
- 7/6/21 (Item 21 from file: 155) 03837227 83032742  
The mechanism of *in situ* reactivation of glycerol-inactivated coenzyme B12-dependent enzymes; glycerol dehydratase and diol dehydratase.
- 7/6/22 (Item 22 from file: 155) 03818946 82119943  
Glycerol fermentation in *Klebsiella pneumoniae*: functions of the coenzyme B12-dependent glycerol and diol dehydratases.
- 7/6/23 (Item 23 from file: 155) 03817061 82099691  
[The molecular basis of manifestation of function for vitamin B12 coenzymes (author's transl)]
- 7/6/24 (Item 24 from file: 155) 03814098 82066743  
Reactive sulphydryl groups of coenzyme B12-dependent diol dehydratase: differential modification of essential and nonessential ones.
- 7/6/25 (Item 25 from file: 155) 03810110 82023979  
Purification and subunit characterization of propanediol dehydratase, a membrane-associated enzyme.
- 7/6/26 (Item 26 from file: 155) 03790172 81085020  
Coenzyme B12-dependent diol dehydratase: chemical modification with 2,3-butanedione and phenylglyoxal.
- 7/6/27 (Item 27 from file: 155) 03783669 81006730  
In situ reactivation of glycerol-inactivated coenzyme B12-dependent enzymes, glycerol dehydratase and diol dehydratase.
- 7/6/28 (Item 28 from file: 155) 03782638 80264192  
Inactivation of dioldehydratase in the presence of a coenzyme-B12 analog.
- 7/6/29 (Item 29 from file: 155) 03775514 80159971  
The synthesis of several immobilized derivatives of vitamin B12 coenzyme and their use as affinity adsorbents for a study of interactions of diol dehydratase with the coenzyme.
- 7/6/30 (Item 30 from file: 155) 03775503 80159893  
Distribution of coenzyme B12-dependent diol dehydratase and glycerol dehydratase in selected genera of Enterobacteriaceae and Protonobacteriaceae.
- 7/6/31 (Item 31 from file: 155) 03260264 79231445  
Stereospecificity and mechanism of adenosylcobalamin-dependent diol dehydratase. Catalysis and inactivation with meso- and dl-2,3-butanediols as substrates.
- 7/6/32 (Item 32 from file: 155) 03115235 79124674  
Role of peripheral side chains of vitamin B12 coenzymes in the reaction catalyzed by dioldehydratase.
- 7/6/33 (Item 33 from file: 155) 03108208 78242158  
Coenzyme B12-dependent diol dehydratase: regulation of apoenzyme synthesis in *Klebsiella pneumoniae* (Aerobacter aerogenes) ATCC 8724.
- 7/6/34 (Item 34 from file: 155) 02985254 77225263  
Immunochemical evidence for the difference between coenzyme-B12-dependent diol dehydratase and glycerol dehydratase.
- 7/6/35 (Item 35 from file: 155) 02963505 80000580  
Resolution of the coenzyme B12-dependent dehydratases of *Klebsiella* sp. and *Citrobacter freundii*.
- 7/6/36 (Item 36 from file: 155) 02963490 80000417  
Hydrogen transfer in catalysis by adenosylcobalamin-dependent diol dehydratase.
- 7/6/37 (Item 37 from file: 155) 02959767 79216215  
Fermentation of 1,2-propanediol with 1,2-ethanediol by some genera of Enterobacteriaceae, involving coenzyme B12-dependent diol dehydratase.
- 7/6/38 (Item 38 from file: 155) 02966899 79186157  
Coenzyme B12-dependent diol dehydratase: purification, subunit heterogeneity, and reversible association.
- 7/6/39 (Item 39 from file: 155) 02908486 77134713  
Mechanism of action of adenosylcobalamin: glycerol and other substrate analogues as substrates and inactivators for propanediol dehydratase—kinetics, stereospecificity, and mechanism.
- 7/6/40 (Item 40 from file: 155) 02907797 77118572  
Studies on the mechanism of the adenosylcobalamin-dependent diol dehydratase reaction by the use of analogs of the coenzyme.
- 7/6/41 (Item 41 from file: 155) 02313552 75146954  
Preparation, properties and biological activities of succinyl derivatives of vitamin B12.
- 7/6/42 (Item 42 from file: 155) 02143923 76184142  
Substrate specificity of coenzyme B12-dependent diol dehydratase: glycerol as both a good substrate and a potent inactivator.
- 7/6/43 (Item 43 from file: 155) 02058873 76039896  
Immobilized diol dehydratase and its use in studies of cobalamin binding and subunit interaction.
- 7/6/44 (Item 44 from file: 155) 02002592 75146949  
Ethanolamine ammonia-lyase: inactivation of the holoenzyme by N2O and the mechanism of action of Coenzyme B12.
- 7/6/45 (Item 45 from file: 155) 01430076 75008121  
Coenzyme B12 dependent diol dehydratase system: Dissociation of the enzyme into two different protein components and some properties of the components.
- 7/6/46 (Item 46 from file: 155) 01317779 74031427  
Activation of diol dehydratase by formamidineum or guanidinium ion, polyatomic monovalent cations having sp<sup>2</sup> nitrogen.
- 7/6/47 (Item 47 from file: 155) 01276156 73196460  
Dissociation of diol dehydratase into two different protein components.
- 7/6/48 (Item 48 from file: 155) 01201378 73047392  
Coenzyme B12 -dependent propanediol dehydratase systems. Ternary complex between apoenzyme, coenzyme, and substrate analog.

7/6/49 (Item 49 from file: 155) 0158359 72238147  
Coenzyme B 12 dependent propanediol dehydratase system. Nature of cobalamin binding and some properties of apoenzyme-coenzyme B 12 analog complexes.

7/6/50 (Item 50 from file: 155) 00961699 72040213  
Propanediol dehydratase system. Role of monovalent cations in binding of vitamin B 12 coenzyme or its analogs to apoenzyme.

7/6/51 (Item 51 from file: 155) 00495878 70000235  
Ternary complex formation of 1,2-propanediol dehydratase, cobamide coenzyme and substrate analogue.

7/6/52 (Item 52 from file: 155) 00727137 68011874  
Coenzyme activity of 5-chlorocobalamin (10-Cl-DBCC) in propanediol dehydratase system.

7/6/53 (Item 53 from file: 155) 00143034 67173019  
[On the mechanism of the propanediol dehydratase reaction] Zum Mechanismus der Propandioldehydratase-Reaktion.

7/6/54 (Item 54 from file: 155) 00102610 67052680  
[On the stereochemistry of the propanediol dehydratase reaction] Zur Stereochemie der Propandioldehydratase-Reaktion.

7/6/55 (Item 1 from file: 5) 13011721 BIOSIS Number: 99011721  
Carbon and electron flow in Clostridium butyricum grown in chemostat culture on glycerol and on glucose Print Number: Biological Abstracts Vol. 102 Iss. 001 Ref. 011721

7/6/56 (Item 2 from file: 5) 11049541 BIOSIS Number: 97249541  
Diol dehydratase and glycerol dehydratase, coenzyme B-12-dependent isozymes Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082721

7/6/57 (Item 3 from file: 5) 11049540 BIOSIS Number: 97249540  
Diol dehydratase from Clostridium glycolicum: The non-B-12-dependent enzyme Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082720

7/6/58 (Item 4 from file: 5) 11049533 BIOSIS Number: 97249533  
Metal ions in Biological Systems. Vol. 30. Metalloenzymes involving amino acid-residue and related radicals Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082713

7/6/59 (Item 5 from file: 5) 5667450 BIOSIS Number: 94072450  
ENZYMES INVOLVED IN ANAEROBIC POLYETHYLENE GLYCOL DEGRADATION BY PELOBACTER-VENETIANUS AND BACTEROIDES STRAIN PG1

7/6/60 (Item 6 from file: 5) 5816885 BIOSIS Number: 83079192  
SOLUBILIZATION OF A MEMBRANE-BOUND DIOL DEHYDRATASE WITH RETENTION OF EPR G EQUALS 2.02 SIGNAL BY USING 2-N CYCLOHEXYLAMINOETHANESULFONIC ACID BUFFER

7/6/61 (Item 7 from file: 5) 5447256 BIOSIS Number: 82092059  
CHARACTERIZATION OF THE ENZYME INVOLVED IN FORMATION OF 2 BUTANOL FROM MESO-2,3 BUTANEDIOL BY LACTIC-ACID BACTERIA

7/6/62 (Item 8 from file: 5) 5150605 BIOSIS Number: 31039920  
SOLUBILIZATION OF MEMBRANE-BOUND AND OXYGEN SENSITIVE ENZYMES WITH 2-N CYCLOHEXYLAMINOETHANESULFONIC-ACID

7/6/63 (Item 9 from file: 5) 5104752 BIOSIS Number: 30117059  
SOLUBILIZATION OF DIOL DEHYDRATASE FROM CLOSTRIDIUM-GLYCOLICUM

7/6/64 (Item 10 from file: 5) 4792137 BIOSIS Number: 79034452  
COENZYMIC FUNCTION OF 1 SUBSTITUTED OR N-6 SUBSTITUTED ANALOGS OF ADENOSYL-COBALAMIN IN THE DIOL DEHYDRATASE EC-4.2.1.28 REACTION

7/6/65 (Item 11 from file: 5) 4650225 BIOSIS Number: 29007540  
DIOL DEHYDRATASE AND GLYCOL METABOLISM IN CLOSTRIDIUM-GLYCOLICUM

7/6/66 (Item 12 from file: 5) 4440303 BIOSIS Number: 78014126  
LIGAND EXCHANGE REACTIONS OF DIOL DEHYDRASE EC-4.2.1.28 BOUND COBALAMINS AND THE EFFECT OF THE NUCLEOSIDE BINDING

7/6/67 (Item 13 from file: 5) 4071485 BIOSIS Number: 76021337  
DIOL DEHYDRATASE EC-4.2.1.28 N TERMINAL AMINO-ACID SEQUENCES AND SUBUNIT STOICHIOMETRY

7/6/68 (Item 14 from file: 5) 4027710 BIOSIS Number: 75075069  
THE MECHANISM OF IN-SITU REACTIVATION OF GLYCEROL INACTIVATED COENZYME B-12 DEPENDENT ENZYMES GLYCEROL DEHYDRATASE EC-4.2.1.30 AND DIOL DEHYDRATASE EC-4.2.1.28

7/6/69 (Item 15 from file: 5) 3664154 BIOSIS Number: 73056521  
REACTIVE SULFHYDRYL GROUPS OF COENZYME B-12 DEPENDENT DIOL DEHYDRASE EC-4.2.1.28 DIFFERENTIAL MODIFICATION OF ESSENTIAL AND NONESSENTIAL ONES

7/6/70 (Item 16 from file: 5) 3642292 BIOSIS Number: 73034659  
PURIFICATION AND SUBUNIT CHARACTERIZATION OF PROPANEDIOL DEHYDRATASE EC-4.2.1.28 A MEMBRANE ASSOCIATED ENZYME

7/6/71 (Item 17 from file: 5) 3318561 BIOSIS Number: 71040860  
COENZYME B-12 DEPENDENT DIOL DEHYDRASE EC-4.2.1.28 CHEMICAL MODIFICATION WITH 2,3 BUTANEDIONE AND PHENYL GLYOXAL

7/6/72 (Item 18 from file: 5) 3237789 BIOSIS Number: 21030192  
STRUCTURE FUNCTION RELATIONSHIP OF VITAMIN B-12 COENZYME ADENOSYL COBALAMIN IN THE DIOL DEHYDRASE EC-4.2.1.28 SYSTEM

7/6/73 (Item 19 from file: 5) 3086882 BIOSIS Number: 70036789  
THE SYNTHESIS OF SEVERAL IMMOBILIZED DERIVATIVES OF VITAMIN B-12 COENZYME AND THEIR USE AS AFFINITY ADSORBENTS FOR A STUDY OF INTERACTIONS OF DIOL DEHYDRASE EC-4.2.1.28 WITH THE COENZYME

7/6/74 (Item 20 from file: 5) 2974674 BIOSIS Number: 69012081  
FERMENTATION OF 1,2 PROPANEDIOL AND 1,2 ETHANEDIOL BY SOME GENERA OF ENTEROBACTERIACEAE INVOLVING COENZYME B-12 DEPENDENT DIOL DEHYDRATASE EC-4.2.1.28

7/6/75 (Item 21 from file: 5) 2801475 BIOSIS Number: 68055382  
COENZYME B-12 DEPENDENT DIOL DEHYDRASE EC-4.2.1.28 PURIFICATION SUBUNIT HETEROGENEITY AND REVERSIBLE ASSOCIATION

7/6/76 (Item 22 from file: 5) 2786880 BIOSIS Number: 68043587  
STEREOSPECIFICITY AND MECHANISM OF ADENOSYL COBALAMIN DEPENDENT DIOL DEHYDRATASE CATALYSIS AND INACTIVATION WITH MESO-2,3 BUTANEDIOL AND RACEMIC 2,3 BUTANEDIOL AS SUBSTRATES

7/6/77 (Item 23 from file: 5) 2756523 BIOSIS Number: 68011430  
ROLE OF PERIPHERAL SIDE CHAINS OF VITAMIN B-12 COENZYMES IN THE REACTION CATALYZED BY DIOL DEHYDRASE EC-4.2.1.28

7/6/78 (Item 24 from file: 5) 2684235 BIOSIS Number: 67021638  
METABOLISM OF 1,2 PROPANEDIOL BY METHANOL UTILIZING BACTERIA AND SOME PROPERTIES OF 1,2 PROPANEDIOL DEHYDROGENATING ENZYME

7/6/79 (Item 25 from file: 5) 2526149 BIOSIS Number: 66073054  
COENZYME B-12 DEPENDENT DIOL DEHYDRATASE EC-4.2.1.28 REGULATION OF APOENZYME SYNTHESIS IN KLEBSIELLA-PNEUMONIAE AEROBACTER-AEROGES ATCC-8724

7/6/80 (Item 26 from file: 5) 2501151 BIOSIS Number: 66048056  
MECHANISM OF ACTION OF ADENOSYL COBALAMIN HYDROGEN TRANSFER IN THE INACTIVATION OF DIOL DEHYDRATASE EC-4.2.1.28 BY GLYCEROL

7/6/81 (Item 27 from file: 5) 2377808 BIOSIS Number: 66004216  
IMMUNOCHEMICAL EVIDENCE FOR THE DIFFERENCE BETWEEN COENZYME B-12 DEPENDENT DIOL DEHYDRATASE EC-4.2.1.28 AND GLYCEROL DEHYDRATASE EC-4.2.1.30

7/6/82 (Item 28 from file: 5) 2183678 BIOSIS Number: 64010598  
MECHANISM OF ACTION OF ADENOSYL COBALAMIN GLYCEROL AND OTHER SUBSTRATE ANALOGS AS SUBSTRATES AND INACTIVATORS FOR PROPANEDIOL DEHYDRATASE EC-4.2.1.28 KINETICS STEREOSPECIFICITY AND MECHANISM

7/6/83 (Item 29 from file: 5) 2166587 BIOSIS Number: 63071007  
STUDIES ON THE MECHANISM OF THE ADENOSYL COBALAMIN DEPENDENT DIOL DEHYDRASE EC-4.2.1.28 REACTION BY THE USE OF ANALOGS OF THE COENZYME

7/6/84 (Item 30 from file: 5) 1721539 BIOSIS Number: 60066107  
A PHYSICAL EXPLANATION OF THE EPR SPECTRUM OBSERVED DURING CATALYSIS BY ENZYMES UTILIZING COENZYME B-12

7/6/85 (Item 31 from file: 5) 1677514 BIOSIS Number: 60022082  
ETHANOL AMINE AMMONIA LYASE INACTIVATION OF THE HOLO ENZYME BY NITROGEN OXIDE AND THE MECHANISM OF ACTION OF COENZYME B-12



- 7/6/86** (Item 32 from file: 5) 1671435 BIOSIS Number: 60016003  
RELATIVE ENANTIOMER BINDING AND REACTION RATES WITH PROPANEDIOL DEHYDRASE EC-4.2.2.28
- 7/6/87** (Item 33 from file: 5) 1083898 BIOSIS Number: 55013830  
FORMATION OF 5-DEOXYADENOSYL DERIVATES OF COBALAMIN C LACTAM AND COBALAMIN C LACTONE BY PROPIONIBACTERIUM SHERMANII IN-VIVO AND IN-VITRO
- 7/6/88** (Item 1 from file: 73) 10002807 EMBASE No: 96181477  
Evidence for enantiomeric-antitopic group discrimination in diol dehydratase-catalyzed dehydration of meso-2,3-butanediol
- 7/6/89** (Item 2 from file: 73) 9133324 EMBASE No: 94072716  
The synthesis of a pyridyl an analog of adenosylcobalamin and its coenzymic function in the diol dehydratase reaction
- 7/6/90** (Item 3 from file: 73) 8280211 EMBASE No: 91302965  
Roles of the D-ribose and 5,6-dimethylbenzimidazole moieties of the nucleotide loop of adenosylcobalamin in manifestation of coenzymic function in the diol dehydratase reaction
- 7/6/91** (Item 4 from file: 73) 7247646 EMBASE No: 88247524  
Acceleration of cleavage of the carbon-cobalt bond of sterically hindered alkylcobalamins by binding to apoprotein of diol dehydratase
- 7/6/92** (Item 5 from file: 73) 6031502 EMBASE No: 86026562  
The binding site for the adenosyl group of coenzyme B<sub>12</sub> in diol dehydratase
- 7/6/93** (Item 6 from file: 73) 5754272 EMBASE No: 84249938  
Propanediol-1,2-dehydratase and metabolism of glycerol of *Lactobacillus brevis*
- 7/6/94** (Item 7 from file: 73) 5710823 EMBASE No: 84206489  
Coenzymic function of 1- or N-substituted analogs of adenosylcobalamin in the diol dehydratase reaction
- 7/6/95** (Item 8 from file: 73) 5125653 EMBASE No: 82130576  
Glycerol fermentation in *Klebsiella pneumoniae*: Functions of the coenzyme B<sub>12</sub> sub 2-dependent glycerol and diol dehydratases
- 7/6/96** (Item 9 from file: 73) 2260075 EMBASE No: 81031200  
In situ reactivation of glycerol-inactivated coenzyme B<sub>12</sub> sub 2-dependent enzymes: glycerol dehydratase and diol dehydratase
- 7/6/97** (Item 10 from file: 73) 1232941 EMBASE No: 79000296  
Coenzyme B<sub>12</sub> sub 2-dependent diol dehydratase: regulation of apoenzymesynthesis in *Klebsiella pneumoniae* (Aerobacter aerogenes) ATCC 8724
- 7/6/98** (Item 11 from file: 73) 949780 EMBASE No: 78117969  
Immunochemical evidence for the difference between coenzyme B<sub>12</sub> sub 2 dependent diol dehydratase and glycerol dehydratase
- 7/6/99** (Item 12 from file: 73) 616302 EMBASE No: 76203083  
Mechanism of action of adenosylcobalamin: 3-fluoro-1,2-propanediol as substrate for propanediol dehydratase. Mechanistic implications
- 7/6/100** (Item 13 from file: 73) 556098 EMBASE No: 76140982  
Coenzyme action of adenosyl 13 epicobalamin in the diol dehydratase system
- 7/6/101** (Item 14 from file: 73) 537094 EMBASE No: 76121511  
A physical explanation of the EPR spectrum observed during catalysis by enzymes utilizing coenzyme B<sub>12</sub> sub 2
- 7/6/102** (Item 15 from file: 73) 427247 EMBASE No: 76007141  
Relative enantiomer binding and reaction rates with propanediol dehydratase
- 7/6/103** (Item 16 from file: 73) 326271 EMBASE No: 75119035  
Coenzyme B<sub>12</sub> sub 2 dependent diol dehydratase system. Dissociation of the enzyme into two different protein components and some properties of the components
- 7/6/104** (Item 1 from file: 351) 011021737 WPI Acc No: 96-518683/199651  
Fermentative production of 1,3-propane-diol useful for polymer production - from carbon substrates using mixed culture of glycerol-producing and diol-producing organisms
- 7/6/105** (Item 2 from file: 351) 011021733 WPI Acc No: 96-518683/199651  
Osmid control *Klebsiella pneumoniae* gene for diol dehydratase - and related transformed microorganisms able to convert glycerol to 1,3-propanediol for polymer production
- 7/7/13** (Item 3 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
08791004 96394290  
Cloning, sequencing, and high level expression of the genes encoding adenosylcobalamin-dependent glycerol dehydratase of *Klebsiella pneumoniae*.
- 7/7/14** (Item 4 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
08790962 96422012  
Cloning, sequencing, and overexpression of the genes encoding coenzyme B<sub>12</sub>-dependent glycerol dehydratase of *Citrobacter freundii*.  
Seyfried M; Daniel R; Gottschalk G  
Institut für Mikrobiologie der Georg-August-Universität, Göttingen, Germany,  
J Bacteriol (UNITED STATES) Oct 1996; 178 (19) p5793-6, ISSN 0021-9193 Journal Code: HH3 Languages: ENGLISH Document type: JOURNAL ARTICLE  
The genes encoding coenzyme B<sub>12</sub>-dependent glycerol dehydratase of *Citrobacter freundii* were cloned and overexpressed in *Escherichia coli*. The B<sub>12</sub>-free enzyme was purified to homogeneity. It consists of three types of subunits whose N-terminal sequences are in accordance with those deduced from the open reading frames dhaB, dhaC, and dhaE, coding for subunits of 60 433 (alpha), 21 487 (beta), and 16 121 (gamma) Da, respectively. The enzyme complex has the composition alpha2beta2gamma2. Amino acid alignments with the subunits of the recently sequenced diol dehydratase of *Klebsiella oxytoca* (T. Tobimatsu, T. Hara, M. Sakaguchi, Y. Kishimoto, Y. Wada, M. Isoda, T. Sakai, and T. Toraya, J. Biol. Chem. 270:7142-7148, 1995) revealed identities between 51.8 and 70.9%.
- 7/7/15** (Item 5 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
08213743 95221362  
Molecular cloning, sequencing, and expression of the genes encoding adenosylcobalamin-dependent diol dehydratase of *Klebsiella oxytoca*.  
Tobimatsu T; Hara T; Sakaguchi M; Kishimoto Y; Wada Y; Isoda M; Sakai T; Toraya T  
Department of Biotechnology, Faculty of Engineering, Okayama University, Japan.  
J Biol Chem (UNITED STATES) Mar 31 1995; 270 (13) p7142-8, ISSN 0021-9258 Journal Code: HHV Languages: ENGLISH Document type: JOURNAL ARTICLE  
The pdd genes encoding adenosylcobalamin-dependent diol dehydratase of *Klebsiella oxytoca* were cloned by using a synthetic oligodeoxynucleotide as a hybridization probe followed by measuring the enzyme activity of each clone. Five clones of *Escherichia coli* exhibited diol dehydratase activity. At least one of them was shown to express diol dehydratase genes under control of their own promoter. Sequence analysis of the DNA fragments found in common in the inserts of these five clones and the flanking regions revealed four open reading frames separated by 10-18 base pairs. The sequential three open reading frames from the second to the fourth (pddA, pddB, and pddC genes) encoded polypeptides of 554, 224, and 173 amino acid residues with predicted molecular weights of 60 348 (alpha), 24 113 (beta), and 19 173 (gamma), respectively. Overexpression of these three genes in *E. coli* produced more than 50-fold higher level of functional apodiol dehydratase than that in *K. oxytoca*. The recombinant enzyme was indistinguishable from the wild-type one of *K. oxytoca* by the criteria of polyacrylamide gel electrophoretic and immunochemical properties. It was thus concluded that these three gene products are the subunits of functional diol dehydratase. Comparisons of the deduced amino acid sequences of the three subunits with other proteins failed to reveal any apparent homology.
- 7/7/11** (Item 11 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
06130099 86129441  
Diol metabolism and diol dehydratase in *Clostridium glycolicum*.  
Hartmanis MG; Siedman TC  
Arch Biochem Biophys (UNITED STATES) Feb 15 1986; 245 (1) p144-52, ISSN 0003-9861 Journal Code: 6SK Languages: ENGLISH Document type: JOURNAL ARTICLE

Levels of the five enzymes involved in the fermentation of 1,2-ethanediol and 1,2-propanediol in the strictly anaerobic bacterium, *Clostridium glycolicum*, were investigated. All enzymes with the exception of the first enzyme in the pathway, diol dehydratase, were found to be constitutive, stable to exposure to oxygen, and present in the cytosol. Diol dehydratase was found to be extremely oxygen sensitive and strongly associated with the cell membrane. Treatment with ionic and nonionic detergents, butanol, phospholipase A2, or osmotic shock procedures failed to solubilize any diol dehydratase activity. Limited proteolysis using subtilisin released small amounts of activity. Diol dehydratase was found to be specific for 1,2-ethanediol and 1,2-propanediol and required the addition of a reducing agent for maximal activity. The enzyme was strongly inhibited by low concentrations of EDTA, ethylene glycol bis(beta-aminoethyl ether)-N,N',N'',N'''-tetraacetic acid or phenanthroline, hydroxylamine, hydroxyurea, and sulfinyl reagents. Addition of adenosylcobalamin or high levels of intrinsic factor did not affect the reaction rate. Irradiation with light also did not inhibit the enzyme activity. These results suggest that the catalytic mechanism of diol dehydratase from *C. glycolicum* does not involve a cobamide coenzyme.

**717122** (Item 22 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All rts. reserv.  
03818946 82119943

Glycerol fermentation in *Klebsiella pneumoniae*: functions of the coenzyme B12-dependent glycerol and diol dehydratases.  
Forage RG; Foster MA  
J Bacteriol (UNITED STATES) Feb 1982; 149 (2) p413-9, ISSN 0021-9193 Journal Code: HH3 Languages: ENGLISH  
Document type: JOURNAL ARTICLE

Glycerol and diol dehydratases are inducible, coenzyme B12-dependent enzymes found together in *Klebsiella pneumoniae* ATCC 25955 during anaerobic growth on glycerol. Mutants of this strain isolated by a novel procedure were separately constitutive for either dehydratase, showing the structural genes for the two enzymes to be under independent control *in vivo*. Glycerol dehydratase and a trimethylene glycol dehydrogenase were implicated as members of a pleiotropic control system that includes glycerol dehydrogenase and dihydroxyacetone kinase for the anaerobic dissimilation of glycerol (the "dia system"). The dehydratase and dehydrogenases were induced by dihydroxyacetone and were jointly constitutive in mutants isolated as constitutive for either the dia system or glycerol dehydratase. These data and the stimulation of growth by Co2+ suggested that glycerol dehydratase and trimethylene glycol dehydrogenase are obligatory enzymes for anaerobic growth on glycerol as the sole carbon source.

**717156** (Item 2 from file: 5) DIALOG(R)File 5 BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
11049541 BIOSIS Number: 97249541

Diol dehydratase and glycerol dehydratase, coenzyme B-12-dependent isozymes

Toraya T  
Dep. Biotechnol., Fac. Eng., Okayama Univ., 3-1-1 Tsushima-Naka, Okayama 700, JAP 0 (0). 1994, 217-254, Full Journal Title: Sigel, H. and A. Sigel (Ed.), Metal Ions in Biological Systems, Vol. 30. Metalloenzymes involving amino acid-residue and related radicals. xxxv+494p.  
Marcel Dekker, Inc.: New York, New York, USA; Basel, Switzerland. ISBN 0-8247-9093-6. ISSN: 0161-5149 Language: ENGLISH Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082721

**717157** (Item 3 from file: 5) DIALOG(R)File 5 BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
11049540 BIOSIS Number: 97249540

Diol dehydratase from *Clostridium glycolicum*: The non-B-12-dependent enzyme

Hartmanis M GN

Kabi Pharmacia BioSci. Cent., Strandbergsgatan 49, S-11287 Stockholm, SWE 0 (0). 1994, 201-215.  
Full Journal Title: Sigel, H. and A. Sigel (Ed.), Metal Ions in Biological Systems, Vol. 30. Metalloenzymes involving amino acid-residue and related radicals. xxxv+494p. Marcel Dekker, Inc.: New York, New York, USA; Basel, Switzerland. ISBN 0-8247-9093-6. ISSN: 0161-5149 Language: ENGLISH Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082720

**717158** (Item 4 from file: 5) DIALOG(R)File 5 BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
11049533 BIOSIS Number: 97249533

Metal Ions in Biological Systems, Vol. 30. Metalloenzymes involving amino acid-residue and related radicals

Sigel H; Sigel A  
Inst. Inorg. Chem., Univ. Basel, CH-4056 Basel, SWI 0 (0). 1994, XXXV+494P. Full Journal Title: Sigel, H. and A. Sigel (Ed.), Metal Ions in Biological Systems, Vol. 30. Metalloenzymes involving amino acid-residue and related radicals. xxxv+494p. Marcel Dekker, Inc.: New York, New York, USA; Basel, Switzerland. ISBN 0-8247-9093-6. ISSN: 0161-5149 Language: ENGLISH Print Number: Biological Abstracts/RRM Vol. 046 Iss. 006 Ref. 082713

This book contains 13 papers discussing metalloenzymes involving amino acid-residue and related radicals. Some of the topics covered include free radical sites and their locations, mechanistic considerations, and enzymes that depend on the metals manganese, iron, cobalt, and copper. The work will be useful for researchers and students in chemistry, biochemistry, biophysics, enzymology, molecular biology, etc. Graphs, diagrams, tables, and charts illustrate the text.

**717159** (Item 5 from file: 5) DIALOG(R)File 5 BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
9567450 BIOSIS Number: 94072450

ENZYMES INVOLVED IN ANAEROBIC POLYETHYLENE GLYCOL DEGRADATION BY PELOBACTER-VENETIANUS AND BACTEROIDES STRAIN PG1  
FRINGS J; SOHRAMME E; SCHINK B  
FAKULTAET FUER BIOLOGIE DER UNIVERSITAET KONSTANZ, POSTFACH 5560, D-7750 KONSTANZ, GERMANY.  
APPL ENVIRON MICROBIOL 58 (7). 1992. 2164-2167. CODEN: AEMID Full Journal Title: Applied and Environmental Microbiology Language: ENGLISH

In extracts of polyethylene glycol (PEG)-grown cells of the strictly anaerobically fermenting bacterium *Pelobacter venetianus*, two different enzyme activities were detected, a diol dehydratase and a PEG-degrading enzyme which was characterized as a PEG acetaldehyde lyase. Both enzymes were oxygen sensitive and depended on a reductant, such as titanium citrate or sulfinyl compounds, for optimal activity. The diol dehydratase was inhibited by various corinoids (adenosylcobalamin, cyanocobalamin, hydroxocobalamin, and methylcobalamin) by up to 37% at a concentration of 100 .mu.M. Changes in ionic strength and the K+ ion concentration had only limited effects on this enzyme activity; glycerol inhibited the enzyme by 95%. The PEG-degrading enzyme activity was stimulated by the same corinoids by up to 80%, inhibited optimal activity in 0.75 M potassium phosphate buffer or in the presence of 4 M KCl, and was only slightly affected by glycerol. Both enzymes were located in the cytoplasmic space. Also, another PEG-degrading bacterium, *Bacteroides strain PG1*, contained a PEG acetaldehyde lyase activity analogous to the corresponding enzyme of *P. venetianus* but no diol dehydratase. Our results confirm that corinoid-influenced PEG degradation analogous to a diol dehydratase reaction is a common strategy among several different strictly anaerobic PEG-degrading bacteria.

**717174** (Item 20 from file: 5) DIALOG(R)File 5 BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
2974674 BIOSIS Number: 69012081

FERMENTATION OF 1,2-PROPANEDIOL AND 1,2-ETHANEDIOL BY SOME GENERA OF ENTEROBACTERIACEAE INVOLVING COENZYME B-12 DEPENDENT DIOL DEHYDRATASE EC-4.2.1.28

TORAYA T; HONDA S; FUKUI S

LAB. IND. BIO-CHEM., DEP. IND. CHEM., FAC. ENG., KYOTO UNIV., SAKYO, KYOTO 606, JPN.  
J BACTERIOL 139 (1). 1979, 39-47. CODEN: JOBAA Full Journal Title: Journal of Bacteriology Language: ENGLISH  
*Klebsiella pneumoniae* (Aerobacter aerogenes) ATCC 8724 grew anaerobically on 1,2-propanediol and 1,2-ethanediol as C and energy sources. Whole cells of the bacterium grown anaerobically on 1,2-propanediol or on glycerol catalyzed conversion of 1,2-diols and aldehydes on the corresponding acids and alcohols. Glucose-grown cells also converted aldehydes, but not 1,2-diols, to acids and alcohols. The presence of activities of coenzyme B12-dependent diol dehydratase, alcohol dehydrogenase, Co-A-dependent aldehyde dehydrogenase, phosphotransacetylase and acetate kinase was demonstrated with crude extracts of 1,2-propanediol-grown cells. The dependence of the levels of these enzymes on growth substrates, together with cofactor requirements in *in vitro* conversion of these substrates, indicates that 1,2-diols are fermented to the corresponding acids and alcohols via aldehydes, acyl-CoA and acyl phosphates. This metabolic pathway for 1,2-diol fermentation was also suggested in some other genera of *Enterobacteriaceae* which grew anaerobically on 1,2-propanediol. When the bacteria were cultivated in a 1,2-propanediol medium not supplemented with cobalt ion, the coenzyme B12-dependent conversion of 1,2-diols to aldehydes was the rate-limiting step in this fermentation. This was because the intracellular concentration of coenzyme B12 was very low in the cells grown in cobalt-deficient medium, since the apoprotein of diol dehydratase was markedly induced in the cells grown in the 1,2-propanediol medium. Better cell yields were obtained when the bacteria were grown anaerobically on 1,2-propanediol. Aerobically grown cells evidently have a different metabolic pathway for utilizing 1,2-propanediol.

**717105** (Item 2 from file: 351) DIALOG(R)File 351:DERWENT WPI (c)1997 Derwent Info Ltd. All rts. reserv.  
011021733 WPI Acc No: 96-518883/199651

Cosmid contg. *Klebsiella pneumoniae* gene for diol dehydratase - and related transformed microorganisms able to convert glycerol to 1,3-propanediol for polymer prodn

Patent Assignee: DUPONT DE NEMOURS & CO E I (DUPO )

Inventor: NAGARAJAN V; NAKAMURA C E

Number of Countries: 061 Number of Patents: 003

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Main IPC	Week
WO 9635795	A1	1996/1114	WO 96/US 6163	A	1996/0502	C12N-015/60	199651 B
AU 9657229	A	1996/1129	AU 9657229	A	1996/0502	C12N-015/60	199712
US 5633362	A	1997/0627	US 95440377	A	1996/0512	C07H-021/02	199727

Priority Applications (No Type Date): US 95440377 A 19960512

Cited Patents: 9, journal ref.

Patent Details:

Patent Kind Lan Pg Filing Notes Application Patent

WO 9635795 A1 E 48

Designated States (National): AL AU BB BG BR CA CN CZ EE GE HU IS JP KP KR LK LR LT LV MG MK MN MX NO NZ PL RO SG SI SK TR TT UA UZ VN

Designated States (Regional): AT BE CH DE DK EA ES FI FR GB GR IE IT KE LS LU MC MW NL OA PT SD SE SZ UG



AU 9637229 A Based on WO 9635795  
US 5633362 A 18  
Abstract (Basic): WO 9635795 A  
Cosmid (A) comprises a DNA fragment (I) of about 35 kb from *Klebsiella pneumoniae* that encodes an active diol dehydratase enzyme (II).  
USE - Cells transformed with (I) or (A) can convert glycerol to 1,3-propanediol (IV) which is a monomer potentially useful for prodn. of polyester fibre, polyurethanes and cyclic cpds.  
ADVANTAGE - This method provides efficient, cost effective and environmentally acceptable prodn. of (IV).  
Dwg. 0/4  
Abstract (Equivalent): US 5633362 A  
A cosmid comprising a DNA fragment of about 35 kb isolated from *Klebsiella pneumoniae* wherein said fragment encodes an active diol dehydratase enzyme having the restriction digest in FIG. 5, columns numbered 4, said cosmid contained within a transformed *E. coli* deposited with the American Type Culture Collection under accession number ATCC 69790. (Fig 5 not suitable for reproduction) Dwg. 0/0  
Derwent Class: A41; D16; E17; F01  
International Patent Class (Main): C07H-02/102; C12N-01/56/0  
International Patent Class (Additional): C07H-02/104; C12N-00/119; C12N-00/121; C12N-00/904; C12N-00/988; C12N-01/56/3; C12N-01/57/4; C12N-01/57/9; C12P-00/718  
96/1 (Item 1 from file: 155) 09265995 97457194  
Glycerol conversion to 1,3-propanediol by *Clostridium pasteurianum* cbing and expression of the gene encoding 1,3-propanediol dehydrogenase.  
96/2 (Item 2 from file: 155) 09229632 97388589  
Anaerobic pathways of glycerol dissimilation by *Enterobacter agglomerans* CNCM 1210: Imitations and regulations.  
96/3 (Item 3 from file: 155) 08016680 94377734  
Phenotypic diversity of anaerobic glycerol dissimilation shown by seven enterobacterial species.  
96/4 (Item 4 from file: 155) 07313946 93122543  
Growth temperature-dependent activity of glycerol dehydratase in *Escherichia coli* expressing the *Citrobacter freundii* dha regulon.  
96/5 (Item 5 from file: 155) 07070052 92121087  
Sugar-glycerol cofermentations in lactobacilli: the fate of lactate.  
96/6 (Item 6 from file: 155) 06924196 92152855  
1,3-Propanediol production by *Escherichia coli* expressing genes from the *Klebsiella pneumoniae* dha regulon.  
96/7 (Item 7 from file: 155) 06901057 90155202  
Anaerobic growth of *Escherichia coli* on glycerol by importing genes of the dha regulon from *Klebsiella pneumoniae*.  
96/8 (Item 8 from file: 155) 05308385 87194586  
*Klebsiella pneumoniae* 1,3-propanediol:NAD<sup>+</sup>-oxido-reductase.  
96/9 (Item 9 from file: 155) 03838735 83049313  
[Coenzyme properties of adenosylcobalamin analogs with modifications in the purine nucleus of the alpha-legend] Koflermentnyye svoystva analogov adenozilkobalamina s izmenennym purinovym iadrom alfa-Iganda.  
96/10 (Item 10 from file: 155) 03825037 82183110  
[Substrate specificity of adenosylcobalamin-dependent glycerol dehydratase. Interaction with enantiomers of 1,2-propanediol] Substratnaya spetsifichnost' adenozilkobalaminzavisimoi gliserolegidratazy. Vzaimodeystvie s enantiomerami 1,2-propandiola.  
96/11 (Item 11 from file: 155) 03151140 77065983  
[Effect of environmental factors on inactivation of B12-dependent glycerol dehydratase from *Aerobacter aerogenes*] Vlianiye faktorov sredy na inaktivatsiyu B12-zavisimoi gliserolegidratazy iz *Aerobacter aerogenes*  
96/12 (Item 12 from file: 155) 03134432 75174520  
Glycerol dehydratase from *Aerobacter aerogenes*.  
96/13 (Item 13 from file: 155) 02620573 79062639  
[Effect of the structure of the nucleoside ligand of cobalamines on their enzymatic properties in a glycerol dehydratase system] Vlianiye struktury nukleozidnogo lganda kobalaminov na kh koflermentnyye svoystva v sisteme gliserolegidratazy.  
96/14 (Item 14 from file: 155) 02491634 78061052  
[9-Adenyltyalkobalamins as inhibitors of adenosylcobalamin-dependent glycerol dehydratase from *Aerobacter aerogenes*] 9-Adenil'tyalkobalaminy kak inhibitory adenozilkobalamin-zavisimoi gliserolegidratazy iz *Aerobacter aerogenes*.  
96/15 (Item 15 from file: 155) 02449780 77242443

Study of the mechanism of action of adenosylcobalamin-dependent glycerol dehydratase from *Aerobacter aerogenes*. II. The inactivation kinetics of glycerol dehydratase complexes with adenosylcobalamin and its analogs.  
96/16 (Item 16 from file: 155) 02449779 77242442  
Study on the mechanism of action of adenosylcobalamin-dependent glycerol dehydratase from *Aerobacter aerogenes*. I. Role of structural components of adenosylcobalamin in the formation of the active site of glycerol dehydratase.  
96/17 (Item 17 from file: 155) 02079984 76089220  
[Role of monovalent cations in reactions catalyzed by glycerol dehydratase from *Aerobacter aerogenes*]  
96/18 (Item 18 from file: 155) 01807381 74300091  
Determination of glycerol dehydratase activity by the coupled enzymic method.  
96/19 (Item 19 from file: 155) 01802219 74150185  
[Determination of glycerol dehydratase activity by the method of coupled enzyme reactions] Opredeleniye aktivnosti gliserolegidratazy metodom sopriazheniia fermentativnykh reaktsii.  
96/20 (Item 20 from file: 155) 01472942 75134080  
[Study of purine analogs of cobamide coenzyme in a glycerol dehydratase system from *aerobacter aerogenes*] Izucheniye purinovykh analogov kobamidnogo koflamenta v sisteme gliserolegidratazy iz *aerobacter aerogenes*  
96/21 (Item 21 from file: 155) 01424920 74269724  
Allosteric interactions in glycerol dehydratase. Purification of enzyme and effects of positive and negative cooperativity for glycerol.  
96/22 (Item 22 from file: 155) 01363562 74080757  
[Formation of glycerol dehydratase by a culture of *Aerobacter aerogenes*, its partial purification and various properties] Obrazovanie gliserolegidratazy kulturoi *Aerobacter aerogenes*, ee chastiichnaya ocshtishka i nekotoryye svoystva.  
96/23 (Item 23 from file: 155) 01244861 75002999  
[Kinetics of irreversible inactivation of holoenzyme and enzyme-substrate complexes of glycerol dehydratase] Kinetika neobratimoi inaktivatsii koflamenta i fermentatsionnykh kompleksov gliserolegidratazy  
96/24 (Item 24 from file: 155) 01209238 73067771  
[Kinetics of the transformation of 1,2-propanediol to propionic aldehyde, catalyzed by glycerol dehydratase from *Aerobacter aerogenes*] Kinetika prevrashcheniia 1,2-propandiola v propionovyi al'degid, kataliziruemogo gliserolegidratazoi iz *Aerobacter aerogenes*.  
96/25 (Item 25 from file: 155) 01103372 70293158  
Purification and properties of glycerol dehydratase.  
96/26 (Item 26 from file: 155) 01081824 68277312  
Mechanism of action of coenzyme B12-dependent glycerol dehydratase.  
96/27 (Item 27 from file: 155) 00218268 67257076  
Enzymatic determination of vita min B12, coenzyme B12, and other cobamide derivatives in picomole quantities by means of glycerol dehydratase from *Aerobacter aerogenes*.  
96/28 (Item 28 from file: 155) 00136925 67124546  
The properties of glycerol dehydratase isolated from *Aerobacter aerogenes*, and the properties of the apoenzyme subunits.  
96/29 (Item 1 from file: 5) 13582798 BIOSIS Number: 95682798  
Biochemical and molecular characterization of coenzyme B-12-dependent glycerol dehydratase from *Citrobacter freundii* Print Number: Biological Abstracts/RRM Vol. 049 Iss. 007 Ref. 118404  
96/30 (Item 2 from file: 5) 13333745 BIOSIS Number: 98333745  
Physiologic mechanisms involved in accumulation of 3-hydroxypropionaldehyde during fermentation of glycerol by *Enterobacter agglomerans* Print Number: Biological Abstracts Vol. 103 Iss. 003 Ref. 036859  
96/31 (Item 3 from file: 5) 12230210 BIOSIS Number: 98830210  
Glycerol dehydratase activity: The limiting step for 1,3-propanediol production by *Clostridium butyricum* DSM 5431 Print Number: Biological Abstracts Vol. 101 Iss. 012 Ref. 180632  
96/32 (Item 4 from file: 5) 10107492 BIOSIS Number: 95107492  
FERMENTATION OF GLYCEROL TO 1,3-PROPANEDIOL IN CONTINUOUS CULTURES OF *CITROBACTER-FREUNDII*  
96/33 (Item 5 from file: 5) 9107519 BIOSIS Number: 93992519  
SUGAR GLYCEROL COFERMENTATIONS IN *LACTOBACILLI* THE FATE OF LACTATE  
96/34 (Item 6 from file: 5) 7479751 BIOSIS Number: 89130770  
UTILIZATION OF GLYCEROL AS A HYDROGEN ACCEPTOR BY *LACTOBACILLUS-REUTERI* PURIFICATION OF 1,3-PROPANEDIOL NAD OXIDOREDUCTASE

- 9655 (Item 2 from file: 73) 6412604 EMBASE No: 87149266  
Klebsiella pneumoniae 1,3-propanediol NADsup + oxidoreductase
- 9656 (Item 3 from file: 73) 1264966 EMBASE No: 79032619  
Effects of the nucleoside ligands structure of cobalamines on their coenzymic properties in glycerol dehydratase
- 9657 (Item 4 from file: 73) 100051 EMBASE No: 78170429  
Study on the mechanism of action of adenosylcobalamin-dependent glycerol dehydratase from Aerobacter aerogenes. II. The inactivation kinetics of glycerol dehydratase complexes with adenosylcobalamin and its analogs
- 9658 (Item 5 from file: 73) 100050 EMBASE No: 78170428  
Study on the mechanism of action of adenosylcobalamin-dependent glycerol dehydratase from Aerobacter aerogenes. I. Role of structural components of adenosylcobalamin in the formation of the active site of glycerol dehydratase
- 9659 (Item 6 from file: 73) 859479 EMBASE No: 78025357  
Influence of environmental factors on the inactivation of Bsub (sub 2 dependent) glycerol dehydratase from Aerobacter aerogenes
- 9660 (Item 7 from file: 73) 630679 EMBASE No: 77007407  
The role of monovalent cations in reactions catalyzed by glycerol dehydratase from Aerobacter aerogenes
- 9661 (Item 8 from file: 73) 516161 EMBASE No: 93310393  
Response to vasoactive neuropeptides in basilar arteries isolated from stroke-prone spontaneously hypertensive rats
- 9662 (Item 9 from file: 73) 468469 EMBASE No: 76048032  
The interaction of epoglycerol dehydratase from Aerobacter aerogenes with 'apurine' analogs of cobamide coenzyme
- 9663 (Item 10 from file: 73) 444734 EMBASE No: 76025321  
Production of glycerol dehydratase by culture of Aerobacter aerogenes, its partial purification, and some properties
- 9664 (Item 11 from file: 73) 372001 EMBASE No: 75167006  
Investigation of purine analogues of the cobamide coenzyme in the glycerol dehydratase system from Aerobacter aerogenes (Russian)
- 9711 (Item 1 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
09265995 97457194  
Glycerol conversion to 1,3-propanediol by Clostridium pasteurianum: cloning and expression of the gene encoding 1,3-propanediol dehydrogenase.  
Luers F, Seyfried M, Daniel R, Gottschalk G  
Institut für Mikrobiologie der Georg-August-Universität, Göttingen, Germany.  
FEMS Microbiol Lett (NETHERLANDS) Sep 15 1997; 154 (2) p337-45, ISSN 0378-1097 Journal Code: FML Languages: ENGLISH Document type: JOURNAL ARTICLE  
When grown on glycerol as sole carbon and energy source, cell extracts of Clostridium pasteurianum exhibited activities of glycerol dehydrogenase, dihydroxyacetone kinase, glycerol dehydratase and 1,3-propanediol dehydrogenase. The genes encoding the latter two enzymes were cloned by colony hybridization using the cha1 gene of Citrobacter freundii as a heterologous DNA probe and expressed in Escherichia coli. The native molecular mass of 1,3-propanediol dehydrogenase (Dha1) is 440,000 Da. The dha1 gene of C. pasteurianum was subcloned and its nucleotide sequence (1158 bp) was determined. The deduced gene product (41,776 Da) revealed high similarity to DhaT of C. freundii (80.5% identity; 89.6% similarity).
- 9714 (Item 4 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
07313946 93122543  
Growth temperature-dependent activity of glycerol dehydratase in Escherichia coli expressing the Citrobacter freundii dha regulon.  
Daniel R, Gottschalk G  
Institut für Mikrobiologie, Georg-August-Universität, Göttingen, FRG.  
FEMS Microbiol Lett (NETHERLANDS) Dec 15 1992; 79 (1-3) p281-5, ISSN 0378-1097 Journal Code: FML Languages: ENGLISH Document type: JOURNAL ARTICLE  
Using the cosmid pWE15, a genomic library of Citrobacter freundii DNA in Escherichia coli ECL707 was prepared and screened for glycerol utilization. Six out of approximately 3000 clones were positive. One clone, harboring the recombinant cosmid pRD1, expressed glycerol dehydratase in high activity when grown at 28 degrees C but not at 37 degrees C. The growth temperature had little effect on the activity of the other enzymes encoded by the dha regulon. When the glycerol-containing medium was supplemented with cornoids, the recombinant E. coli strain produced 1,3-propanediol in high amounts at 28 degrees C.
- 9715 (Item 5 from file: 155) DIALOG(R)File 155:MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All its. reserv.  
07070352 92121087  
Sugar-glycerol fermentations in lactobacilli: the fate of lactate.
- 9635 (Item 7 from file: 5) 7479748 BIOSIS Number: 89130767  
PURIFICATION AND CHARACTERIZATION OF GLYCEROL DEHYDRATASE FROM LACTOBACILLUS-REUTERI
- 9636 (Item 8 from file: 5) 4521051 BIOSIS Number: 78094874  
ANAEROBIC REDUCTION OF GLYCEROL TO 1,3-PROPANEDIOL BY LACTOBACILLUS-BREVIS AND LACTOBACILLUS-BUCHNERI
- 9637 (Item 9 from file: 5) 4402667 BIOSIS Number: 77077994  
COBALT C-CORRINOIDS THE DERIVATIVES OF VITAMIN B-12 PSEUDOTRANSFORMS AS CORRINOID ENZYME INHIBITORS
- 9638 (Item 10 from file: 5) 4347088 BIOSIS Number: 77022415  
SOME PHYSICO-CHEMICAL FEATURES OF GLYCEROL DEHYDRATASE CATALYZED REACTIONS
- 9639 (Item 11 from file: 5) 4343221 BIOSIS Number: 77018548  
PRODUCTION OF 3-HYDROXY-PROPIONALDEHYDE FROM GLYCEROL
- 9640 (Item 12 from file: 5) 4167203 BIOSIS Number: 26019546  
COENZYME PROPERTIES OF ADENOSYL COBALAMIN ANALOGS WITH A CHANGED PURINE NUCLEUS OF THE ALPHA-LIGAND
- 9641 (Item 13 from file: 5) 4079098 BIOSIS Number: 76028949  
COENZYME PROPERTIES OF ADENOSYL COBALAMIN ANALOGS WITH MODIFICATIONS IN THE ALPHA-LIGAND
- 9642 (Item 14 from file: 5) 3847492 BIOSIS Number: 24054851  
SUBSTRATE SPECIFICITY OF ADENOSYL COBALAMIN DEPENDENT GLYCEROL DEHYDRATASE EC-4.2.1.30 INTERACTION WITH ENANTIOMERS OF 1,2-PROPANEDIOL
- 9643 (Item 15 from file: 5) 3693569 BIOSIS Number: 73085936  
GLYCEROL FERMENTATION IN KLEBSIELLA-PNEUMONIAE FUNCTIONS OF THE COENZYME-B-12 DEPENDENT GLYCEROL AND DIOL DEHYDRATASES
- 9644 (Item 16 from file: 5) 2974535 BIOSIS Number: 68012042  
INTERACTION OF SUBSTRATES AND THEIR ANALOGS WITH ADENOSYL COBALAMIN DEPENDENT GLYCEROL DEHYDRATASE
- 9645 (Item 17 from file: 5) 2944727 BIOSIS Number: 19049636  
PARTICIPATION OF CYCLIC AMP IN REGULATION OF COENZYME B-12 DEPENDENT GLYCEROL DEHYDRATASE EC-4.2.1.30 SYNTHESIS FROM KLEBSIELLA-PNEUMONIAE ATCC-25965
- 9646 (Item 18 from file: 5) 2944720 BIOSIS Number: 19049629  
ADENOSYL COBALAMIN DEPENDENT GLYCEROL DEHYDRATASE EC-4.2.1.30 INTERACTION WITH SUBSTRATES AND THEIR ANALOGS
- 9647 (Item 19 from file: 5) 2937208 BIOSIS Number: 19042117  
ENZYMATIC ESTIMATION OF VITAMIN B-12
- 9648 (Item 20 from file: 5) 2856392 BIOSIS Number: 18028803  
INTERACTION OF SUBSTRATES AND THEIR ANALOGS WITH ADENOSYL COBALAMIN DEPENDENT GLYCEROL DEHYDRATASE EC-4.2.1.30
- 9649 (Item 21 from file: 5) 2835422 BIOSIS Number: 18007833  
EFFECT OF STRUCTURE OF NUCLEOSIDE LIGAND OF COBALAMINS ON THEIR COENZYME PROPERTIES IN THE GLYCEROL DEHYDRATASE EC-4.2.1.30 SYSTEM
- 9650 (Item 22 from file: 5) 2782252 BIOSIS Number: 68037159  
EFFECT OF THE NUCLEOSIDE LIGAND STRUCTURE OF COBALAMINS ON THEIR COENZYME PROPERTIES IN THE GLYCEROL DEHYDRATASE SYSTEM
- 9651 (Item 23 from file: 5) 2775647 BIOSIS Number: 68030554  
SEARCH FOR NEW MEDICINAL PREPARATIONS ON THE BASIS OF VITAMIN B-12 DERIVATIVES SYNTHESIS AND STUDY OF THE PHYSICO-CHEMICAL AND COENZYME PROPERTIES OF ADENOSYL COBALAMIN DERIVATIVES
- 9652 (Item 24 from file: 5) 2106317 BIOSIS Number: 63010737  
THE ROLE OF E-PROPANAMIDE GROUP OR THE CORRIN MACRO CYCLE IN THE MANIFESTATION OF COENZYMIC PROPERTIES OF THE COBALAMIDE COENZYME
- 9653 (Item 25 from file: 5) 1666115 BIOSIS Number: 60010683  
STUDY OF PURINE ANALOGS OF THE COBALAMIDE COENZYME IN THE GLYCEROL DEHYDRATASE SYSTEM FROM AEROBACTER-AEROGES
- 9654 (Item 1 from file: 73) 8406923 EMBASE No: 92083103  
Sugar-glycerol fermentations in lactobacilli: The fate of lactate

- Veiga da Cunha M; Foster MA  
Department of Biochemistry, University of Oxford, United Kingdom  
J Bacteriol (UNITED STATES) Feb 1992; 174 (3) p1013-9. ISSN 0021-9193 Journal Code: HH3 Languages: ENGLISH  
Document type: JOURNAL ARTICLE
- The simultaneous fermentation of glycerol and sugar by *Lactobacillus brevis* B22 and *Lactobacillus buchneri* B190 increases both the growth rate and total growth. The reduction of glycerol to 1,3-propanediol by the lactobacilli was found to influence the metabolism of the sugar cofermented by channeling some of the intermediate metabolites (e.g., pyruvate) towards NADH-producing (rather than NADH-consuming) reactions. Ultimately, the absolute requirement for NADH to prevent the accumulation of 3-hydroxypropionaldehyde leads to a novel lactate-glycerol cofermentation, as a result, additional ATP can be made not only by (i) converting pyruvate to acetate via acetyl phosphate rather than to the ethanol usually found and (ii) oxidizing part of the intermediate pyruvate to acetate instead of the usual reduction to lactate but also by (iii) reoxidation of accumulated lactate to acetate via pyruvate. The conversion of lactate to pyruvate is probably catalyzed by NAD-independent lactate dehydrogenases that are found only in the cultures oxidizing lactate and producing 1,3-propanediol, suggesting a correlation between the expression of these enzymes and a raised intracellular NAD/NADH ratio. The enzymes metabolizing glycerol (glycerol dehydratase and 1,3-propanediol dehydrogenase) were expressed in concert without necessary induction by added glycerol, although their expression may also be influenced by the intracellular NAD/NADH ratio set by the different carbohydrates fermented.
- 9/718 (Item 8 from file: 155) DIALOG(R)File 155: MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All rts. reserv.  
05308385 87194586  
*Klebsiella pneumoniae* 1,3-propanediol NAD+ oxidoreductase.  
Johnson EA, Lin EC  
J Bacteriol (UNITED STATES) May 1987; 169 (5) p2050-4. ISSN 0021-9193 Journal Code: HH3 Contract/Grant No.: 5-R01-GM11983, GM, NIGMS Languages: ENGLISH Document type: JOURNAL ARTICLE
- Fermentative utilization of glycerol, a more reduced carbohydrate than aldoses and ketoses, requires the disposal of the two extra hydrogen atoms. This is accomplished by sacrificing an equal quantity of glycerol via an auxiliary pathway initiated by glycerol dehydratase. The product, 3-hydroxypropionaldehyde, is then reduced by 1,3-propanediol NAD+ oxidoreductase (1,3-propanediol dehydrogenase; EC 1.1.1.202), resulting in the regeneration of NAD+ from NADH. The pathway for the assimilation of glycerol is initiated by an NAD-linked endoenzyme. In *Klebsiella pneumoniae* the two pathways are encoded by the *cha* region which is inducible only anaerobically. In this study 1,3-propanediol:NAD+ oxidoreductase was purified from cells grown anaerobically on glycerol. The enzyme was immunologically distinct from the NAD-linked glycerol dehydrogenase and was an octamer or hexamer of a polypeptide of 45,000 +/- 3,000 daltons. When tested as a dehydrogenase, only 1,3-propanediol served as a substrate; no activity was detected with ethanol, 1-propanol, 1,2-propanediol, glycerol, or 1,4-butanediol. The enzyme was inhibited by chelators of divalent cations. An enzyme preparation inhibited by alpha, alpha'-dipyridyl was reactivated by the addition of Fe2+ or Mn2+ after removal of the chelator by gel filtration. As for glycerol dehydrogenase, 1,3-propanediol oxidoreductase is apparently inactivated by oxidation during aerobic metabolism, under which condition the enzyme becomes superfluous.
- 9/7112 (Item 12 from file: 155) IALOG(R)File 155: MEDLINE(R) (c) format only 1997 Knight-Ridder Info. All rts. reserv.  
03134432 75174520  
Glycerol dehydratase from *Aerobacter aerogenes*.  
Johnson BC, Strojinski A, Schneider Z  
Methods Enzymol (UNITED STATES) 1975; 42 p315-23. ISSN 0076-6879 Journal Code: MVA Languages: ENGLISH  
Document type: JOURNAL ARTICLE
- 9/7129 (Item 1 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
13582798 BIOSIS Number: 95882798  
Biochemical and molecular characterization of coenzyme B-12-dependent glycerol dehydratase from *Citrobacter freundii* Daniel R; Seyfried M; Gottschalk G  
Inst. Microbiol. Georg-August-Universität Göttingen, Grisebachstr. 8, 37077 Göttingen, Germany  
Abstracts of the General Meeting of the American Society for Microbiology 97 (0) 1997: 353. Full Journal Title: 97th General Meeting of the American Society for Microbiology, Miami Beach, Florida, USA, May 4-8, 1997. Abstracts of the General Meeting of the American Society for Microbiology ISSN: 1060-2011 Language: ENGLISH Print Number: Biological Abstracts/RRM Vol. 049 Iss. 007 Ref. 118404
- 9/7131 (Item 3 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
12230210 BIOSIS Number: 98830210  
Glycerol dehydratase activity. The limiting step for 1,3-propanediol production by *Clostridium butyricum* DSM 5431 Abbad-Andaloussi S; Guedon E; Spieser E; Petitdemange H  
Lab. Chimie Biol. I, Univ. Henri Poincaré Nancy I, BP 239, 54506 Vandœuvre-les-Nancy Cedex, France  
Letters in Applied Microbiology 22 (4) 1996: 311-314. Full Journal Title: Letters in Applied Microbiology ISSN: 0268-8254 Language: ENGLISH Print Number: Biological Abstracts Vol. 101 Iss. 012 Ref. 18652
- Glycerol catabolism by *Clostridium butyricum* DSM 5431 into acetate, butyrate and 1,3-propanediol (1,3-PD) was studied in chemostat culture. The fact that the intracellular concentrations of NADH (18-22 mu-mol g-1 dry cell mass) were extremely high suggested that the dehydratase activity was the rate limiting step in 1,3-PD formation. This limitation was proved by the addition of propionaldehyde, another substrate of propanediol dehydrogenase, into the culture medium. This resulted in an increase in (i) glycerol utilization, (ii) biomass formation and (iii) product biosynthesis.
- 9/7132 (Item 4 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
10107492 BIOSIS Number: 95107492  
FERMENTATION OF GLYCEROL TO 1,3-PROPANEDIOL IN CONTINUOUS CULTURES OF *CITROBACTER FREUNDII* BOENIGK R; BOWEN S; GOTTSCHALK G  
INSTITUT FUER MIKROBIOLOGIE, GEORG-AUGUST-UNIVERSITAET GOETTINGEN, GRISEBACHSTRASSE 8, W-3400 GOETTINGEN, GERMANY.  
APPL MICROBIOL BIOTECHNOL 38 (4) 1993; 453-457. CODEN: AMBID Full Journal Title: Applied Microbiology and Biotechnology Language: ENGLISH
- The conversion of glycerol to 1,3-propanediol by *Citrobacter freundii* DSM 30040 was optimized in single- or two-stage continuous cultures. The productivity of 1,3-propanediol formation was higher under glycerol limitation and increased with the dilution rate (D) to a maximum of 3.7 g cndiol. l-1 cndiol. h-1. Glycerol dehydratase seemed to be the rate-limiting step in 1,3-propanediol formation. Conditions for the two-stage fermentation process were as follows: first stage: glycerol limitation (250 mM), pH 7.2, D = 0.1 h-1, 32 degree C; second stage, additional glycerol, pH 6.6, D = 0.05 h-1, 28 degree C. Under these conditions 876 mM glycerol were consumed, the final 1,3-propanediol concentrations was 1 mM, and the overall productivity, 1.38 g cndiol. l-1 cndiol. h-1.
- 9/7133 (Item 5 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
9107519 BIOSIS Number: 93092519  
SUGAR GLYCEROL COFERMENTATIONS IN LACTOBACILLI THE FATE OF LACTATE  
VEIGA DA CUNHA M; FOSTER MA  
MICROBIOL UNIT, DEP. BIOCHEM. UNIV. OXFORD, OXFORD OX1 3QU, UK.  
J BACTERIOL 174 (3) 1992; 1013-1019. CODEN: JOBAA Full Journal Title: Journal of Bacteriology Language: ENGLISH
- The simultaneous fermentation of glycerol and sugar by *Lactobacillus brevis* B22 and *Lactobacillus buchneri* B190 increases both the growth rate and total growth. The reduction of glycerol to 1,3-propanediol by the lactobacilli was found to influence the metabolism of the sugar cofermented by channeling some of the intermediate metabolites (e.g., pyruvate) towards NADH-producing (rather than NADH-consuming) reactions. Ultimately, the absolute requirement for NADH to prevent the accumulation of 3-hydroxypropionaldehyde leads to a novel lactate-glycerol cofermentation. As a result, additional ATP can be made not only by (i) converting pyruvate to acetate via acetyl phosphate rather than to the ethanol usually found and (ii) oxidizing part of the intermediate pyruvate to acetate instead of the usual reduction to lactate but also by (iii) reoxidation of accumulated lactate to acetate via pyruvate. The conversion of lactate to pyruvate is probably catalyzed by NAD-independent lactate dehydrogenases that are found only in the cultures oxidizing lactate and producing 1,3-propanediol, suggesting a correlation between the expression of these enzymes and a raised intracellular NAD/NADH ratio. The enzymes metabolizing glycerol (glycerol dehydratase and 1,3-propanediol dehydrogenase) were expressed in concert without necessary induction by added glycerol, although their expression may also be influenced by the intracellular NAD/NADH ratio set by the different carbohydrates fermented.
- 9/7135 (Item 7 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
7479748 BIOSIS Number: 89130767  
PURIFICATION AND CHARACTERIZATION OF GLYCEROL DEHYDRATASE FROM *LACTOBACILLUS-REUTERI* TALARICO T L; DOBRGOSZ W J  
DEP. MICROBIOL., NORTH CAROLINA STATE UNIV., RALEIGH, N.C. 27695.  
APPL ENVIRON MICROBIOL 56 (4) 1990; 1195-1197. CODEN: AEMID Full Journal Title: Applied and Environmental Microbiology Language: ENGLISH
- A coenzyme B12-dependent glycerol dehydratase from *Lactobacillus reuteri* has been purified and characterized. The dehydratase has a molecular weight of approximately 200,000, and sodium dodecyl sulfate-polyacrylamide gel electrophoresis yielded a single major band with a molecular weight of 52,000. Km values for substrates and coenzyme B12 were in the millimolar and the submicromolar range, respectively.
- 9/7136 (Item 8 from file: 5) DIALOG(R)File 5: BIOSIS PREVIEWS(R) (c) 1997 BIOSIS. All rts. reserv.  
4521051 BIOSIS Number: 78094874  
ANAEROBIC REDUCTION OF GLYCEROL TO 1,3-PROPANEDIOL BY *LACTOBACILLUS-BREVIS* AND *LACTOBACILLUS-BUCHNERI* SCHUETZ H; RADLER F  
INSTITUT FUER MIKROBIOLOGIE UND WEINFORSCHUNG, UNIVERSITAET MAINZ, POSTFACH 3980, D-6500 MAINZ.  
SVST APPL MICROBIOL 5 (2) 1984; 169-178. CODEN: SAMID Full Journal Title: Systematic and Applied Microbiology Language: ENGLISH

L1 13 S (DIOL OR GLYCEROL) (2N)(DEHYDRASE OR DEHYDRATASE)

8. A isolated gene encoding an active alcohol dehydrogenase comprising a contiguous sequence which consists of SEQ ID NO: 2.

9. A transformed microorganism comprising a host microorganism and the heterologous gene of claim 7 or claim 8.

10. A transformed microorganism comprising E. coli DH5 alpha. and the DNA sequence of claim 7 or claim 8.

US PAT NO: 5,164,309 :IMAGE AVAILABLE: L1: 11 of 13

#### ABSTRACT:

A process of the microbiological preparation of 1,3 propane diol from glycerol in growth media of suitable bacterial strains is described, accompanied by the addition of a cosubstrate in the form of a H-donor and the separation of the propane diol formed. It is characterized in that a biomass is formed in a growth phase from the selected bacterial strain and accompanied by feeding with glycerol and, if necessary, while substantially excluding the H-donor until a stationary growth phase occurs and b) further glycerol and H-donor matched to the biomass are added to the resulting stationary cell suspension for increased 1,3-propane diol formation. This process makes it possible to produce 1,3-propane diol in a high yield from glycerol with a small amount of unobjectionable by-products in a batchwise manner or in continuous form, following immobilization.

#### We claim:

1. In a process for the microbiological preparation of 1,3-propanediol by cultivating in a growth medium containing glycerol and a bacterial strain which is able to convert the glycerol into 1,3-propanediol and isolating the 1,3-propanediol thus obtained, the improvement which comprises the steps of:  
(i) forming a biomass by culturing a bacterial strain from the Citrobacter genus in the growth medium containing glycerol, wherein the formation of the biomass is carried out with the substantial exclusion of any H donor, permitting the bacterial cells to reach a stationary cell phase; thereafter adding to said biomass additional glycerol and a sugar as an H-donor to the biomass, while keeping the cells in essentially a stationary phase; and (iv) then isolating the 1,3-propanediol thus prepared.

2. The process according to claim 1 wherein said strain is a strain of Citrobacter freundii.

3. The process according to claim 1 wherein step (i) is performed under anaerobic conditions.

4. The process according to claim 1 wherein step (ii) is performed under anaerobic conditions.

5. The process according to claim 1 wherein a pH-value of approximately 6.5 to 8.5 is maintained in steps (i) and (ii).

6. The process according to claim 1 wherein steps (i) and (ii) are performed in a mineral medium.

7. The process according to claim 1 wherein step (i) is concluded by the addition of a predetermined quantity of phosphate or nitrogen source.

8. The process according to claim 7 wherein an ammonium salt is used as the nitrogen source or a potassium dihydrogen phosphate is used as the phosphate source.

9. The process according to claim 1 wherein glycerol is initially present in step (iii) in the amount of 0.2 to 1.5 molar concentration.

10. The process according to claim 1 wherein glycerol is initially present in step (i) in approximately 0.1 to 0.4 molar concentration.

11. The process according to claim 1 wherein said biomass obtained in step (i) is immobilized before step (ii).

12. The process according to claim 11 wherein said immobilization is carried out with calcium alginate.

US PAT NO: 4,962,027 :IMAGE AVAILABLE: L1: 12 of 13

#### ABSTRACT:

A method is disclosed for producing 3-hydroxypropionaldehyde (3-HPA) from glycerol by culturing the bacterium Klebsiella pneumoniae having the identifying characteristics of NRRL B-4011, under aerobic conditions, in an aqueous nutrient medium containing glycerol and a compound that causes 3-HPA to be accumulated by blocking the conversion of 3-HPA to trimethylene glycol. This process is particularly useful for the production, from renewable resources, of acrylic acid, an industrially important polymerizable monomer used in the manufacture of synthetic polymers and plastics and which is presently derived from fossil fuel sources.

#### We claim:

1. A method for the production of 3-hydroxypropionaldehyde (3-HPA) from glycerol, which comprises culturing the bacterium Klebsiella pneumoniae NRRL B-4011 or subcultures thereof, under aerobic conditions, in an aqueous nutrient medium containing an amount of glycerol effective for the induction of "glycerol""dehydratase"" and the production of recoverable quantity of 3-HPA, and an amount of semicarbazide hydrochloride sufficient to prevent the conversion of 3-HPA to trimethylene glycol, until a recoverable quantity of 3-HPA is produced.

2. The method of claim 1 wherein said bacterium is first grown in an aqueous nutrient medium containing a carbon source which induces the production of dehydratase enzyme and further incubated in an aqueous medium containing glycerol and semicarbazide hydrochloride.

3. The method of claim 2 wherein said carbon source is glycerol, 1,2-propanediol, or 1,2-ethanediol.

\*\*\*\*\* STN Columbus \*\*\*\*\*

(FILE 'HOME' ENTERED AT 14:56:30 ON 11 DEC 1997)

FILE 'MEDLINE' ENTERED AT 14:56:38 ON 11 DEC 1987

L1 E HYDRO LYASES/CT

L2 2938 S E9

L3 E SACCHAROMYCES/CT

L4 38378 S E3, E4

L5 145 S L1 AND L2

L6 77716 S CLONING, MOLECULAR/CT

L7 14 S L3 AND L4

L5 ANSWER 1 OF 14 MEDLINE

T1 Gene identification using the yeast two-hybrid system.

L5 ANSWER 2 OF 14 MEDLINE

T1 The bifunctional DCOH protein binds to HNF1 independently of its 4-alpha-carbinolamine dehydratase activity.

L5 ANSWER 3 OF 14 MEDLINE

T1 Roles of the FabA and FabZ beta-hydroxyacyl-acyl carrier protein dehydratases in Escherichia coli fatty acid biosynthesis.

L5 ANSWER 4 OF 14 MEDLINE

T1 Mutants that show increased sensitivity to hydrogen peroxide reveal an important role for the pentose phosphate pathway in protection of yeast against oxidative stress.

L5 ANSWER 5 OF 14 MEDLINE

T1 Sticky- and polymerase chain reaction method for systematic gene disruption in Saccharomyces cerevisiae.

L5 ANSWER 6 OF 14 MEDLINE

T1 Cloning of the Candida glabrata TRP1 and HIS3 genes, and construction of their disruptant strains by sequential integrative transformation.

L5 ANSWER 7 OF 14 MEDLINE

T1 Molecular cloning and characterization of the Schizosaccharomyces pombe his3 gene for use as a selectable marker.

L5 ANSWER 8 OF 14 MEDLINE

T1 Cloning of the dihydroxyacid dehydratase-encoding gene (ILDV3) from Saccharomyces cerevisiae.

L5 ANSWER 9 OF 14 MEDLINE

T1 Molecular genetics in Saccharomyces kluyveri: the HIS3 homolog and its use as a selectable marker gene in S. kluyveri and Saccharomyces cerevisiae.

L5 ANSWER 10 OF 14 MEDLINE

T1 Molecular cloning of the imidazoleglycerolphosphate dehydratase gene of Trichoderma harzianum by genetic complementation in Saccharomyces cerevisiae using a direct expression vector.

L5 ANSWER 11 OF 14 MEDLINE

T1 Cloning of deplD, arnD and asd of Leptospira interrogans serovar icterohaemorrhagiae, and nucleotide sequence of the asd gene.

L5 ANSWER 12 OF 14 MEDLINE

T1 Molecular cloning and characterization of the arnD gene encoding 3-dehydroquinase from Salmonella typhi.

L5 ANSWER 13 OF 14 MEDLINE

T1 Characterization of a laaA gene and an ARS element from Mucor circinelloides.

L5 ANSWER 14 OF 14 MEDLINE

T1 Isopropylmalate dehydratase from yeast.

L5 ANSWER 5 OF 14 MEDLINE AN 96437975 MEDLINE

T1 Sticky-end polymerase chain reaction method for systematic gene disruption in Saccharomyces cerevisiae

AU Maifahi M; Gaillardin C; Nicaud J M

CS Institut National Agronomique Paris-Grignon, Laboratoire de Genetique Moleculaire et Cellulaire, INRA CNRS, Thiverval-Grignon, France.

SO YEAST, (1996 Jul) 12 (9) 859-68. Journal code: YEA. ISSN: 0749-503X. CY ENGLAND: United Kingdom DT

Journal Article; (JOURNAL ARTICLE) LA English FS Priority Journals

OS GENBANK:246259 EM 9702 EW 19970204

AB We describe a new procedure for the generation of plasmids containing a large promoter and terminator region of a gene of interest, useful for gene disruption. In a two-step polymerase chain reaction (PCR), a fragment, corresponding to the terminator and promoter regions separated by a 16 bp sequence containing a rare restriction site (e.g. AscI), is synthesized (7-p fragment). This PCR fragment is cloned in vectors presenting a rare blunt-end cloning site and a yeast marker for selection in Saccharomyces cerevisiae (TRP1, HIS3 and KanMX). The final plasmids are directly for gene disruption after linearization



- by the enzyme (e.g. AscI) specific for the rare restriction site. This approach was used to disrupt three open reading frames identified during the sequencing of COS14-1 from chromosome XIV of *S. cerevisiae*.
- CT Check Tags: Support, Non-U.S. Gov't Base Sequence \*\*\*\*Cloning, Molecular: MT, methods\*\*\* Deoxyribonucleases, Type II Site-Specific: ME, metabolism DNA, Fungal ME, metabolism Fungal Proteins: GE, genetics \*Genes, Fungal Genetic Markers Genetic Vectors \*\*\*\*Hydro-Lyases: GE, genetics\*\*\* Kanamycin Resistance: GE, genetics Models, Genetic Molecular Sequence Data \*Mitogenesis \*Polymerase Chain Reaction: MT, methods \*\*\*\*Saccharomyces cerevisiae: GE, genetics\*\*\* Selection (Genetics) Transformation, Genetic
- CN EC 3.1.21- (transdioxynibonuclease AscI); EC 3.1.21.4 (Deoxyribonucleases, Type II Site-Specific); EC 4.2.1. (Hydro-Lyases); EC 4.2.1.19 (imidazoleglycyerophosphate dehydratase); 0 (DNA, Fungal); 0 (Fungal Proteins); 0 (Genetic Markers); 0 (Genetic Vectors); 0 (TRP1 protein)
- L5 ANSWER 6 OF 14 MEDLINE AN 96096521 MEDLINE
- T1 Cloning of the *Candida glabrata* TRP1 and HIS3 genes, and construction of their disruptant strains by sequential integrative transformation.
- AU Kitada K; Yamaguchi E; Aisawa M
- CS Department of Mycology, Nippon Roche Research Center, Kanagawa, Japan.
- SO GENE, (1995 Nov 20) 165 (2) 203-6. Journal code: FOP. ISSN: 0378-1119. CY Netherlands DT Journal; Article; (JOURNAL ARTICLE) LA English FS Priority Journals
- OS GENBANK-U31470 GENBANK-U31471 EM 9603
- AB The *Candida glabrata* (Cg) TRP1 and HIS3 genes have been isolated by complementation of the *Saccharomyces cerevisiae* (Sc) *trp1* and *his3* mutants, respectively. Cg TRP1 encodes a polypeptide of 217 amino acids (aa), whose aa sequence is 58% identical to that of Sc TRP1. Cg HIS3 encodes a polypeptide of 210 aa, whose aa sequence is 73% identical to that of the Sc HIS3. Both Cg TRP1 and HIS3 were disrupted by sequential integrative transformation where the Sc URA3 was used as a selection marker for transformation. The resulting auxotrophic strain of *his3-* and *trp1-* was used to examine the ability of the Sc genes to complement the Cg mutations; Sc HIS3 and TRP1 complemented the Cg *his3-* and *trp1-* mutations, respectively.
- CT Amino Acid Sequence Base Sequence \*Candida: GE, genetics \*\*\*\*Cloning, Molecular\*\*\* \*Fungal Proteins: GE, genetics \*Genes, Structural: Fungal GE, genetics Genetic Complementation Test \*\*\*\*Hydro-Lyases: GE, genetics\*\*\* Molecular Sequence Data \*Mitogenesis Restriction Mapping \*\*\*\*Saccharomyces cerevisiae: GE, genetics\*\*\* Sequence Analysis, DNA Homology, Amino Acid Transformation, Genetic
- CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.19 (imidazoleglycyerophosphatedehydratase); 0 (Fungal Proteins); 0 (TRP1 protein)
- L5 ANSWER 7 OF 14 MEDLINE
- AN 94211206 MEDLINE
- T1 Molecular cloning and characterization of the *Schizosaccharomyces pombe* *his3* gene for use as a selectable marker.
- AU Burke J D; Gould K L
- CS Department of Cell Biology, School of Medicine, Vanderbilt University, Nashville, TN 37232..
- NC GM 47728-01 (NIGMS)
- SO MOLECULAR AND GENERAL GENETICS, (1994 Jan) 242 (2) 169-76. Journal code: NGP. ISSN: 0026-8925. CY GERMANY; Germany, Federal Republic of
- T Journal; Article; (JOURNAL ARTICLE) LA English FS Priority Journals OS GENBANK-L19523; GENBANK-L19524 EM 9407
- AB A DNA fragment which carries the *his3* gene of *Schizosaccharomyces pombe* has been isolated and characterized for use as a selectable marker in transformations. The *his3* gene encodes the imidazole acetyl phosphate transaminase enzyme (E.C.2.6.1.9), which is responsible for converting imidazole acetyl-P to histidinol-P in step 8 of histidine biosynthesis. The nucleotide sequences of a 2196 bp gene fragment and a corresponding cDNA clone were determined. Three intron sequences punctuate the 1451 bp coding region which generates a predicted polypeptide of 384 amino acids with a molecular mass of 42736 daltons. Northern analysis of *his3* mRNAs indicates that the transcript is approximately 1.6 kb in size. Steady-state levels are down-regulated by nitrogen limitation but are unaffected by histidine starvation. The deduced amino acid sequence was compared to the *Saccharomyces cerevisiae* HIS5, *Escherichia coli* HisC, and *Salmonella typhimurium* HisC proteins, all of which are imidazole acetyl phosphate transaminases. The *S. pombe his3* protein was 49.5% identical to the *S. cerevisiae* HIS5 protein and 21.5% identity was found when all four proteins were compared. The shuttle vector pBG1 to the *S. cerevisiae* HIS5 protein smallest functional region of *his3* and the *S. pombe ars1* sequence into pUC18 for use in transformation of *His3-S. pombe* strains. New *S. pombe* strains in which the *his3* gene was deleted have also been constructed.
- CT Check Tags: Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S. Amino Acid Sequence Base Sequence \*\*\*\*Cloning, Molecular\*\*\* DNA, Fungal GE, genetics \*Escherichia coli: GE, genetics \*Genes, Fungal Genetic Markers Histidine, BI, biosynthesis \*\*\*\*Hydro-Lyases: GE, genetics\*\*\* Molecular Sequence Data Restriction Mapping \*\*\*\*Saccharomyces cerevisiae: GE, genetics\*\*\* Salmonella typhimurium: GE, genetics \*\*\*Schizosaccharomyces: GE, genetics
- Schizosaccharomyces: ME, metabolism Sequence Homology, Amino Acid Transcription, Genetic RN 7006-35-1 (Histidine) CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.19 (imidazoleglycyerophosphate dehydratase); 0 (DNA, Fungal); 0 (Genetic Markers) GEN *his3*
- L5 ANSWER 8 OF 14 MEDLINE AN 94131281 MEDLINE
- T1 Cloning of the dihydroxyacid dehydratase-encoding gene (ILV3) from *Saccharomyces cerevisiae*.
- AU Velasco J A; Cansado J; Pena M C; Kawakami T; Laborda J; Notario V
- CS Department of Radiation Medicine, Georgetown University Medical Center, Washington, DC 20007..
- SO GENE, (1993 Dec 31) 137 (2) 179-85. Journal code: FOP. ISSN: 0378-1119. CY Netherlands DT Journal; Article; (JOURNAL ARTICLE) LA English FS Priority Journals
- OS GENBANK-L13975; GENBANK-L11589; GENBANK-L11590; GENBANK-L11591; GENBANK-L11592; GENBANK-L11593; GENBANK-L11594; GENBANK-Z15047; GENBANK-Z15048; GENBANK-L24529 EM 9405
- AB The biosynthesis of branched-chain amino acids (aa) involves three shared pathways through which pyruvate or alpha-ketobutyrate are converted into alpha-keto acids, precursors of valine, leucine or isoleucine. In eukaryotes, few of these common enzymes have been purified to homogeneity, and the whole complement of biosynthetic genes has not been cloned from a single species. In yeasts, most of these genes (ILV genes) have been cloned and sequenced, with the exception of that coding for dihydroxyacid dehydratase (DAD, EC 4.2.1.9), the third enzyme in the common pathways. We have isolated *Saccharomyces cerevisiae* genomic sequences by hybridization to an oligodeoxyribonucleotide (oligo) probe designed from a highly conserved domain among bacterial DAD-encoding genes. The cloned sequences have been located to *S. cerevisiae* chromosome X, mapped within 0.4 centimorgans (cM) of the *ilv3* locus, and found to complement the *ilv3* mutations of various yeast strains. Nucleotide (nt) and aa sequence analyses of the longest open reading frame (ORF) located within the cloned sequences identified them as the ILV3 gene, which codes for the yeast DAD. With our cloning of ILV3, yeast becomes the only eukaryotic system from which all ILV genes have been cloned, thus allowing direct molecular analyses of their regulation.
- L5 ANSWER 9 OF 14 MEDLINE AN 93289813 MEDLINE
- T1 Molecular genetics in *Saccharomyces kluyveri*: the HIS3 homolog and its use as a selectable marker gene in *S. kluyveri* and *Saccharomyces cerevisiae*.
- AU Weinstock K G; Strathern J N
- CS Laboratory of Eukaryotic Gene Expression, ABL-Basic Research Program, NCI-Frederick Cancer Research and Development Center, MD 21702-1201..
- SO YEAST, (1993 Apr) 9 (4) 351-61. Journal code: YEA. ISSN: 0749-503X. CY ENGLAND. United Kingdom DT Journal; Article; (JOURNAL ARTICLE) LA English FS Priority Journals
- OS GENBANK-Z14125 EM 9309
- AB We cloned the *Saccharomyces kluyveri* HIS3 homolog, k-HIS3, and made a partial deletion of the gene. The k-HIS3 gene complemented a HIS3 deletion in *S. cerevisiae*. The DNA sequences of the open reading frames (ORF-s) of the HIS3 homologs are 70% identical at the DNA level and 83% identical at the deduced amino acid level. The ORF upstream of the k-HIS3 gene is related to the PET56 gene of *S. cerevisiae* found upstream of the HIS3 gene of *S. cerevisiae*. The ORF downstream from the k-HIS3 gene is not related to the DED1 gene found downstream of the HIS3 gene in *S. cerevisiae*.
- CT Amino Acid Sequence Base Sequence Chromosome Mapping \*\*\*\*Cloning, Molecular\*\*\* \*Genes, Fungal: GE, genetics Genetic Markers \*\*\*\*Hydro-Lyases: GE, genetics\*\*\* Molecular Sequence Data Mitogenesis \*\*\*\*Saccharomyces cerevisiae: GE, genetics\*\*\* Selection (Genetics) Sequence Analysis, DNA Transformation, Genetic Ureid ME, metabolism RN 66-22-8 (Uracil)
- CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.19 (imidazoleglycyerophosphate dehydratase); 0 (Genetic Markers)
- GEN *HIS3*; PET56, URA3
- L5 ANSWER 10 OF 14 MEDLINE AN 93024323 MEDLINE
- T1 Molecular cloning of the imidazoleglycyerophosphate dehydratase gene of *Trichoderma harzianum* by genetic complementation in *Saccharomyces cerevisiae* using a direct expression vector.
- AU Goldman G H; Dewaele S; Herrera-Estrella A; Geremia R A; Van Montagu M; Contreras R
- CS Laboratorium voor Genetica, Universiteit Gent, Belgium..
- SO MOLECULAR AND GENERAL GENETICS, (1992 Sep) 234 (3) 481-8. Journal code: NGP. ISSN: 0026-8925. CY GERMANY; Germany, Federal Republic of DT Journal; Article; (JOURNAL ARTICLE)
- LA English FS Priority Journals OS GENBANK-Z11528 EM 9301
- AB The *Trichoderma harzianum* imidazoleglycyerophosphate dehydratase gene (*igh*) has been isolated by complementation of a *Saccharomyces cerevisiae his3* mutant using a direct expression vector. This *Escherichia coli*-yeast shuttle vector was developed to allow efficient cloning and expression of cDNA libraries. The cDNA is 627 nucleotides long and codes for a protein of 209 amino acids with an apparent molecular mass of 22,466 daltons. The predicted protein sequence showed 63.6%, 58.7%, and 38.4% identity respectively to the corresponding enzymes from *S. cerevisiae*, *Pichia pastoris* and *E. coli*. Northern analysis showed that the expression of the *igh* gene in *T. harzianum* is not inhibited by external histidine and the level of *igh* mRNA was about threefold higher in cells starved of histidine.
- CT Check Tags: Support, Non-U.S. Gov't Amino Acid Sequence Base Sequence \*\*\*\*Cloning, Molecular\*\*\* Fungal Proteins: GE, genetics Gene Expression \*Genes, Structural: Fungal Genetic Complementation Test Genetic Vectors \*\*\*\*Hydro-Lyases: GE, genetics\*\*\* Molecular Sequence Data RNA, Messenger: GE, genetics \*\*\*Saccharomyces cerevisiae: GE, genetics \*\*\*Trichoderma: GE, genetics
- CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.19 (imidazoleglycyerophosphate dehydratase); 0 (Fungal Proteins); 0 (RNA, Messenger) GEN *igh*
- L5 ANSWER 14 OF 14 MEDLINE
- AN 89200982 MEDLINE
- T1 Isopropylmalate dehydratase from yeast.
- AU Kohlhaw G B

SO METHODS IN ENZYMOLOGY, (1988) 166 423-9. Journal code: MVA, ISSN: 0076-6879, CY United States DT  
Journal: Article, (JOURNAL ARTICLE) LA English FS Priority Journals EM 8907  
CT Chromatography, Affinity: MT, methods \*\*\* Cloning, Molecular \*\*\* Enzyme Stability \*\*\* Hydro-Lyases: GE, genetics\*\*\*  
\*\*\*\*Hydro-Lyases: IP, isolation & purification\*\*\* \*\*\*\*Hydro-Lyases: ME, metabolism\*\*\* Indicators and Reagents Kinetics  
\*\*\*\*Saccharomycetes cerevisiae: EN, enzymology\*\*\* \*\*\*\*Saccharomycetes cerevisiae: GE, genetics\*\*\* Spectrophotometry, Ultraviolet: MT, methods  
CN EC 4.2.1. (Hydro-Lyases); EC 4.2.1.33 (3-isopropylmalate dehydratase); 0 (Indicators and Reagents)

\*\*\*\*\*  
\*  
\* WELCOME TO MESSENGER (APS TEXT) AT USPTO \*  
\*  
\*  
\* THE USPTO PRODUCTION FILES ARE CURRENT THROUGH: \*  
\* JUNE 9 1998 FOR U.S. PATENT TEXT DATA. \*  
\* JUNE 9 1998 FOR U.S. CURRENT CLASSIFICATION DATA. \*  
\* JUNE 9 1998 FOR U.S. PATENT IMAGE DATA. \*  
\*\*\*\*\*  
\* WELCOME TO THE \*  
\* U.S. PATENT TEXT FILE \*  
\*\*\*\*\*

(FILE "USPAT" ENTERED AT 14:39:16 ON 16 JUN 1998)  
L1 13 S (DIOL OR GLYCEROL)(ZN)(DEHYDRASE OR DEHYDRATASE)  
L2 3 S DHAT  
L3 31 S DHAB?

L2  
1. 5,686,276, NOV. 11, 1997, BIOCONVERSION OF A FERMENTABLE CARBON SOURCE TO 1,3-PROPANEDIOL BY A SINGLE MICROORGANISM; LISA ANNE LAFFEND, ET AL., 435/158, 252.31, 252.33 [IMAGE AVAILABLE]  
2. 5,086,386, FEB. 4, 1992, METHOD AND APPARATUS FOR BENCHMARKING THE WORKING SET OF WINDOW-BASED COMPUTER SYSTEMS; NAYEEM ISLAM, 707/202, 364/264, 264.3, 280, 280.6, 281.3, 282, 285, 286, 286.3, 927.2, 927.4, 927.63, 927.81, 928, 929.12, 931, 931.5, 932, 932.1, 932.4, 932.5, 946.2, 950, 950.3, 950.4, 957, 957.1, 957.8, 962, 962.4, 975.4, DIG.1, DIG.2; 395/182.14 [IMAGE AVAILABLE]  
3. 3,948,331, APR. 6, 1976, TRACK ASSEMBLY FOR SNOWMOBILES; RICHARD E. ESCH, 305/132, 180/193 [IMAGE AVAILABLE]

US PAT NO: 5,686,276 [IMAGE AVAILABLE] L2: 1 OF 3  
SUMMARY: BSUM(14) IN KLEBSIELLA PNEUMONIAE AND CITROBACTER FREUNDII, THE GENES ENCODING THE FUNCTIONALLY LINKED ACTIVITIES OF GLYCEROL DEHYDRATASE (DHAB), 1,3-PROPANEDIOL OXIDOREDUCTASE ("DHAT"), GLYCEROL DEHYDROGENASE (DHAD), AND DIHYDROXYACETONE KINASE (DHAK) ARE ENCOMPASSED BY THE DHAB REGULON. THE DHAB REGULONS FROM CITROBACTER AND KLEBSIELLA.

DETDESC DETD(60) THE... ACHIEVED BY PLACING THE NECESSARY STRUCTURAL GENES UNDER THE CONTROL OF ALTERNATE PROMOTORS AS HAS BEEN DEMONSTRATED FOR 1,3-PROPANEDIOL OXIDOREDUCTASE ("DHAT") FROM C. FREUNDII AND DIOL DEHYDRATASE FROM K. OXYTOCA ATCC 8724 (DANIEL ET AL., J. BACTERIOL. 177, 2161 (1995) AND...

L3  
1. 5,753,723, MAY 19, 1998, DENTURE FIXATIVE WITH AN ADHESION PROMOTER; TIANG SHING CHANG, ET AL., 523/120, 106/35, 514/574, 524/42, 239, 321, 549, 559 [IMAGE AVAILABLE]  
2. 5,750,591, MAY 12, 1998, DENTURE ADHESIVE CONTAINING PARTIAL IRONUM, CALCIUM, SODIUM GANTREZ SALT; HAL C. CLARKE, ET AL., 523/120, 433/228.1; 523/118; 524/45, 559, 525/670 [IMAGE AVAILABLE]  
3. 5,723,106, MAR. 3, 1998, REDUCED ALCOHOL MOUTHWASH ANTISEPTIC AND ANTISEPTIC PREPARATION; R. MICHAEL BUCH, ET AL., 424/49, 58 [IMAGE AVAILABLE]  
4. 5,699,269, DEC. 16, 1997, METHOD FOR PREDICTING CHEMICAL OR PHYSICAL PROPERTIES OF CRUDE OILS; TERRENCE RODNEY ASHE, ET AL., 702/30, 436/29, 60 [IMAGE AVAILABLE]

5. 5,686,181, DEC. 9, 1997, DENTURE FIXATIVE; TIANG-SHING CHANG, ET AL., 523/118, 430/180, 523/120, 524/28, 45, 55, 377, 439, 440 [IMAGE AVAILABLE]  
6. 5,686,276, NOV. 11, 1997, BIOCONVERSION OF A FERMENTABLE CARBON SOURCE TO 1,3-PROPANEDIOL BY A SINGLE MICROORGANISM; LISA ANNE LAFFEND, ET AL., 435/158, 252.31, 252.33 [IMAGE AVAILABLE]  
7. 5,650,479, JUL. 22, 1997, INTERFACIALLY POLYMERIZED POLYESTER FILMS; PAUL G. GLUGLA, ET AL., 528/194, 95/43, 54; 210/500.21, 500.26, 528/176, 193 [IMAGE AVAILABLE]  
8. 5,569,581, OCT. 29, 1996, ALTERATION AND PREDICTION OF MALE FERTILITY USING SEMINAL PLASMA AND ITS COMPONENTS; GARY KILLIAN, ET AL., 435/4; 424/520, 435/806 [IMAGE AVAILABLE]  
9. 5,561,177, OCT. 1, 1996, HYDROCARBON FREE DENTURE ADHESIVE; NILOFAR KHALEDI, ET AL., 524/35, 433/180, 523/120, 524/43, 45, 313, 492 [IMAGE AVAILABLE]  
10. 5,543,443, AUG. 6, 1996, DENTURE STABILIZING COMPOSITIONS; JAYANTH RAJAIAH, ET AL., 523/120, 522/148, 523/116, 118, 524/28, 31, 45, 55, 261, 267, 377, 522, 557, 525/100, 101, 102, 207, 328.9, 366, 474, 477, 478, 479, 526/279, 528/15, 26, 31, 32, 33, 374 [IMAGE AVAILABLE]  
11. 5,461,155, OCT. 24, 1995, ORGANIC SOLUBLE METAL-AZO AND METAL-AZOMETHINE DYES; TERRANCE P. SMITH, ET AL., 546/12 [IMAGE AVAILABLE]  
12. 5,424,058, JUN. 13, 1995, DENTURE STABILIZING COMPOSITIONS COMPRISING A MIXED PARTIAL SALT OF A LOWER ALKYL VINYL ETHER-MALEIC ACID COPOLYMER; JAYANTH RAJAIAH, ET AL., 424/49, 106/35, 523/120, 525/328.9, 366, 370, 526/240 [IMAGE AVAILABLE]  
13. 5,405,836, APR. 11, 1995, PET FOODS WITH WATER-SOLUBLE ZINC COMPOUND COATING FOR CONTROLLING MALODOROUS BREATH; THOMAS RICHAR, ET AL., 514/23, 424/49, 53, 439, 442, 426/72, 74, 805 [IMAGE AVAILABLE]  
14. 5,314,998, MAY 24, 1994, ORGANIC SOLVENT-SOLUBLE METAL-AZO AND METAL-AZOMETHINE DYES; TERRANCE P. SMITH, ET AL., 534/701, 710, 711, 713, 723 [IMAGE AVAILABLE]  
15. 5,304,616, APR. 19, 1994, DENTURE STABILIZING COMPOSITIONS HAVING IMPROVED HOLD; JAYANTH RAJAIAH, ET AL., 526/240, 523/118, 120, 525/327.8 [IMAGE AVAILABLE]  
16. 5,242,834, SEP. 7, 1993, ANALYSIS OF ALUMINUM IN AMINO ACIDS BY HIGH PERFORMANCE LIQUID CHROMATOGRAPHY; DURGA V. SUBRAMANIAN, 439/73, 73/61.52, 210/656, 436/74, 161, 174, 175, 182 [IMAGE AVAILABLE]  
17. 5,225,514, JUL. 6, 1993, AZO CONTAINING POLYURETHANES FOR DRUG DELIVERY TO THE LARGE INTESTINES; YOSHIMARU KIMURA, ET AL., 528/76, 514/772.3, 528/85 [IMAGE AVAILABLE]  
18. 5,165,914, NOV. 24, 1992, ORAL COMPOSITIONS CONTAINING ZINC LACTATE COMPLEXES; RICHARD S. VLOCK, 424/62, 49, 641, 642, 643, 673, 676 [IMAGE AVAILABLE]  
19. 5,094,845, MAR. 10, 1992, ORAL COMPOSITIONS CONTAINING ZINC GLUCONATE COMPLEXES; RICHARD S. VLOCK, 424/62, 49, 53, 55, 613, 641, 643, 673 [IMAGE AVAILABLE]  
20. 5,073,604, DEC. 17, 1991, DENTURE STABILIZING COMPOSITIONS; KENNETH T. HOLEVA, ET AL., 525/327.8; 523/120, 525/327.9, 328.9, 366, 370, 526/240 [IMAGE AVAILABLE]  
21. 5,050,692, SEP. 24, 1991, METHOD FOR DIRECTIONAL DRILLING OF SUBTERRANEAN WELLS; HERBERT W. BEINGRABEN, 175/61, 74, 76, 256 [IMAGE AVAILABLE]  
22. 4,980,391, DEC. 25, 1990, DENTURE ADHESIVES AND METHODS FOR PREPARING SAME; LORI D. KUMAR, ET AL., 524/45, 106/35, 523/120, 524/492 [IMAGE AVAILABLE]  
23. 4,948,580, AUG. 14, 1990, MUCO-BIOADHESIVE COMPOSITION; IVAN BROWNING, 514/772.5, 424/434, 435, 443, 447, 448, 484, 514/944, 969 [IMAGE AVAILABLE]  
24. 4,937,066, JUN. 26, 1990, ZINC CONTAINING ORAL COMPOSITIONS; RICHARD S. VLOCK, 424/62, 49, 53, 55, 613, 614, 641, 643, 673 [IMAGE AVAILABLE]

SO PCT Int. Appl., 109 pp. CODEN: PIXXD2  
PI WO 9635795 A1 961114  
DS W: AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG

AI WO 96-US6705 960510 PRAI US 95-440293 950512 DT Patent LA English  
AB A process is provided for the bioconversion of a carbon substrate, preferably glucose, to 1,3-propanediol by a single organism utilizing microorganisms contg. the genes encoding for an active glycerol or diol dehydratase enzyme. Specifically, the glycerol dehydratase gene of *Klebsiella pneumoniae* is used to prep. a transgenic microorganism capable of forming 1,3-propanediol from glucose in high yield. A cosmid covering the dha regulon of *K. pneumoniae* was cloned and the gene for the dehydratase (dhaB1, dhaB2, dhaB3) and the propanediol dehydrogenase were cloned and expressed in a variety of prokaryotic and eukaryotic microbial hosts with the manu. of the propanediol from glucose or maltose demonstrated.

L9 ANSWER 2 OF 3 CAPLUS COPYRIGHT 1998 ACS  
AN 1997-6102 CAPLUS DN 126:30403  
TI Process for making 1,3-propanediol from carbohydrates using mixed microbial cultures  
IN Haynie, Sharon Loretta; Wagner, Lorraine Winona  
PA E.I. Du Pont De Nemours and Company, USA; Haynie, Sharon Loretta; Wagner, Lorraine Winona  
SO PCT Int. Appl., 30 pp. CODEN: PIXXD2  
PI WO 9635799 A1 961114  
DS W: AL, AU, BB, BG, BR, CA, CN, CZ, EE, GE, HU, IS, JP, KP, KR, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG

AI WO 96-US6161 960502 PRAI US 95-440379 950512 DT Patent LA English  
AB The present invention provides a process for the biotransformation of a carbohydrate C source to 1,3-propanediol using mixed yeast and bacterial cultures wherein the carbohydrate is 1st Fermented to glycerol by the yeast cell and then converted to 1,3-propanediol by the bacterial cell contg. an active diol or glycerol dehydratase enzyme. In this process both the yeast and bacterial cultures are supported on the same C source and 1,3-propanediol is isolated from the media.

L9 ANSWER 3 OF 3 CAPLUS COPYRIGHT 1998 ACS  
AN 1995-841557 CAPLUS DN 124:46914  
TI Rapid expansion of the physical and genetic map of the chromosome of *Clostridium perfringens* CPN50  
AU Katayama, Sei-Ichi; Dupuy, Bruno; Garnier, Thierry; Cole, Stewart T.  
CS Unite Genetique Moleculaire Bacterienne, Inst. Pasteur, Paris, 75724, Fr.  
SO J. Bacteriol. (1995), 177(19), 5680-5 CODEN: JOBAAY; ISSN: 0021-9193 DT Journal LA English  
AB The phys. map of the 3.6-megabase chromosome of *Clostridium perfringens* CPN50 was extended by positioning sites for the endonucleases SfiI and I-CeuI, and in parallel, the gene map was expanded by using a genome scanning strategy. This involved the cloning and sequencing of random chromosomal fragments, identification of the functions of the putative genes by database searches, and then hybridization anal. The current gene map comprises almost 100 markers, many of which encode housekeeping functions while others are involved in sporulation or pathogenesis. Strikingly, most of the virulence genes were found to be confined to a 1200-kb segment of the chromosome near oriC, while the pleiotropic regulatory locus, virRS, was situated toward the putative replication terminus. A comparison of the gene maps of 3 endospore-forming bacilli, *C. perfringens*, *Clostridium beijerinckii*, and *Bacillus subtilis*, revealed a similar order and distribution of key sporulation and heat shock genes which might reflect an ancient evolutionary relationship.

L13 ANSWER 1 OF 8 CAPLUS COPYRIGHT 1998 ACS  
TI Metabolic engineering of propanediol pathways  
L13 ANSWER 2 OF 8 CAPLUS COPYRIGHT 1998 ACS  
TI Metabolic engineering of an improved 1,3-propanediol fermentation (*Klebsiella pneumoniae*, "Bacillus" *licheniformis*)  
L13 ANSWER 3 OF 8 CAPLUS COPYRIGHT 1998 ACS  
TI Production of 1,3-propanediol from glycerol by recombinant bacteria expressing recombinant diol dehydratase  
L13 ANSWER 4 OF 8 CAPLUS COPYRIGHT 1998 ACS  
TI Bioconversion of a fermentable carbon source to 1,3-propanediol by a single microorganism expressing a foreign glycerol or diol dehydratase gene  
L13 ANSWER 5 OF 8 CAPLUS COPYRIGHT 1998 ACS  
TI Process for making 1,3-propanediol from carbohydrates using mixed microbial cultures  
L13 ANSWER 6 OF 8 CAPLUS COPYRIGHT 1998 ACS

25. 4,817,740, APR. 4, 1989, APPARATUS FOR DIRECTIONAL DRILLING OF SUBTERRANEAN WELLS; HERBERT W. BEIMGABEN, 175/74, 76, 256 [IMAGE AVAILABLE]  
26. 4,747,415, MAY 31, 1988, METHOD AND DEVICE FOR MEASURING PENILE RIGIDITY; PIERRE LAVOISIER, 600/687, 507 [IMAGE AVAILABLE]  
27. 4,717,260, JAN. 5, 1988, TIME DIFFERENTIAL CORRECTING ANALOG TIMEPIECE OF TWENTY-FOUR HOUR SYSTEM; SHIGERU TSUJI, 368/21, 968/167, DIG. 1 [IMAGE AVAILABLE]  
28. 4,560,013, DEC. 24, 1985, APPARATUS FOR DIRECTIONAL DRILLING AND THE LIKE OF SUBTERRANEAN WELLS; HERBERT W. BEIMGABEN, 175/73, 325.2 [IMAGE AVAILABLE]  
29. 4,404,088, SEP. 13, 1983, THREE-STAGE HYDROCRACKING PROCESS; ROBERT W. BACHTEL, ET AL., 208/59, 111 [IMAGE AVAILABLE]  
30. 3,926,577, DEC. 16, 1975, CORROSION INHIBITOR FOR VANADIUM-CONTAINING FUELS; MICHAEL J. ZETLMEISL, ET AL., 44/320, 354; 252/387 [IMAGE AVAILABLE]  
31. 3,691,408, SEP. 12, 1972, METHOD AND MEANS FOR THERMOELECTRIC GENERATION OF ELECTRICAL ENERGY; JOHN B. ROSSO, 310/306; 62/5; 136/209, 211 [IMAGE AVAILABLE]  
US PAT. NO.: 5,686,276 [IMAGE AVAILABLE] L3: 6 OF 31  
SUMMARY: BSJW(14) IN *KLEBSIELLA PNEUMONIAE* AND *CITROBACTER FREUNDII*, THE GENES ENCODING THE FUNCTIONALLY LINKED ACTIVITIES OF GLYCEROL DEHYDRATASE ("DHA8"), 1,3-PROPANEDIOL OXIDOREDUCTASE (DHAT), GLYCEROL DEHYDROGENASE (DHAD), AND DIHYDROXYACETONE KINASE (DHAK) ARE ENCOMPASSED BY THE DHA REGULON. THE DHA REGULONS FROM . . .  
\*\*\*\*\* Welcome to STN International \*\*\*\*\*  
\*\*\*\*\* STN Columbus \*\*\*\*\*  
(FILE 'HOME' ENTERED AT 16:35:10 ON 15 JUN 1998)  
FILE 'REGISTRY' ENTERED AT 16:35:25 ON 15 JUN 1998  
L1 40618 S 1, 3-PROPANEDIOL  
L2 7000 S GLYCEROL  
L3 74 S DIHYDROXYACETONE  
FILE 'CAPLUS' ENTERED AT 16:36:41 ON 15 JUN 1998  
FILE 'REGISTRY' ENTERED AT 16:44:28 ON 15 JUN 1998  
L4 1 S GLYCEROL DEHYDRATASE  
FILE 'CAPLUS' ENTERED AT 16:44:53 ON 15 JUN 1998  
L5 61 S L4  
L6 94627 S ASPERGILLUS OR SACCHAROMYCES OR ZYGOSACCHAROMYCES OR PICHIA OR KLUYVEROMYCES OR CANDIDA OR HANSENULA  
L7 136502 S DEBARYOMYCES OR MUCOR OR TORULOPSIS OR METHYLOBACTER OR SALMONELLA OR BACILLUS OR STREPTOMYCES OR PSEUDOMONAS  
L8 222558 S L6 OR L7  
L9 3 S L5 AND L8  
L10 6439 S 1, 3-PROPANEDIOL  
L11 108 S L8 AND L10 NOT L9  
L12 219 S 604-63-2P/IT  
L13 8 S L12 AND L8  
L9 ANSWER 1 OF 3 CAPLUS COPYRIGHT 1998 ACS  
AN 1997-34085 CAPLUS DN 126:58953  
TI Bioconversion of a fermentable carbon source to 1,3-propanediol by a single microorganism expressing a foreign glycerol or diol dehydratase gene  
IN Laffard, Lisa Anne; Nagarajan, Vasantha; Nakamura, Charles Edwin  
PA E.I. Du Pont De Nemours and Company, USA; Genencor International, Inc.; Laffard, Lisa Anne; Nagarajan, Vasantha; Nakamura, Charles Edwin



- TI Microbial production and downstream processing of 2,3-butanediol
- L13 ANSWER 7 OF 8 CAPLUS COPYRIGHT 1998 ACS
- TI Fermentative manufacture of 1,3-propanediol from glycerol
- L13 ANSWER 8 OF 8 CAPLUS COPYRIGHT 1998 ACS
- TI Neutral solvent production from halophilic, photolithotrophically grown algae by linked fermentations
- L13 ANSWER 1 OF 8 CAPLUS COPYRIGHT 1998 ACS
- AN 1998:56526 CAPLUS DN 128-87891
- TI Metabolic engineering of propanediol pathways
- AU Cameron, D. C.; Altaras, N. E.; Hoffman, M. L.; Shaw, A. J.
- CS Department of Chemical Engineering, University of Wisconsin/Madison, Madison, WI, USA
- SO Biotechnol. Prog. (1998), 14(1), 116-125 CODEN: BIPRET; ISSN: 8756-7938 PB American Chemical Society DT Journal, General Review LA English
- AB A review with many refs. Microbial ferment. is an important technol. for the conversion of renewable resources to chems. In this paper, the authors describe the application of metabolic engineering for the development of two new ferment. processes: the microbial conversion of sugars to 1,3-propanediol (1,3-PD) and 1,2-propanediol (1,2-PD). A variety of naturally occurring organisms ferment glycerol to 1,3-PD, but no natural organisms ferment sugars directly to 1,3-PD. The authors first describe the fed-batch ferment. Of glycerol to 1,3-PD by *Klebsiella pneumoniae*. They then present various approaches for the conversion of sugars to 1,3-PD, including mixed-culture ferment., cofeimentation of glycerol and glucose, and metabolic engineering of a "sugars to 1,3-PD" pathway in a single organism. Results are reported for the expression of genes from the *K. pneumoniae* 1,3-PD pathway in "Saccharomyces" cerevisiae. The best naturally occurring organism for the ferment. of sugars to 1,2-PD is *Thermoanaerobaculum thermoaccharolyticum*. The authors describe the ferment. of several different sugars to 1,2-PD by this organism in batch and continuous culture. They report that *Escherichia coli* strains engineered to express either alcoholose reductase or glycerol dehydrogenase convert glucose to (R)-1,2-PD. The authors then analyze the ultimate potential of ferment. Processes for the prodn. of propanediols. Linear optimization studies indicate that, under aerobic conditions, propanediol yields that approach the theor. max. are possible and CO<sub>2</sub> is the primary coproduct. Without the need to produce acetate, final product titers in the range of 100 g/L should be possible; the high titers and low coproduct levels should make product recovery and purifn. straightforward. The examples given in this paper illustrate the importance of metabolic engineering for ferment. process development in general.
- L13 ANSWER 2 OF 8 CAPLUS COPYRIGHT 1998 ACS
- AN 1997:517535 CAPLUS DN 127:123805
- TI Metabolic engineering of an improved 1,3-propanediol fermentation (*Klebsiella pneumoniae*, "Bacillus" licheniformis)
- AU Skraly, Frank Anthony
- CS Univ. of Wisconsin, Madison, WI, USA
- SO (1997) 221 pp. Avail.: UMI, Order No. DA9716075 From: Diss. Abstr. Int., B 1997, 58(3), 1414 DT Dissertation LA English AB Unavailable
- L13 ANSWER 7 OF 8 CAPLUS COPYRIGHT 1998 ACS
- AN 1990:234037 CAPLUS DN 112:234037
- TI Fermentative manufacture of 1,3-propanediol from glycerol
- IN Kretschmann, Josef; Carduck, Franz Josef; Deckwer, Wolf Dieter; Tag, Carmen
- PA Henkel K.-G.a.A., Fed. Rep. Ger.; Gesellschaft fuer Biotechnologische Forschung m.b.H. (GBF)
- SO Ger. Offen., 7 pp. CODEN: GWXXBX
- PI DE 3829618 A1 900315 AI DE 88-3829618 880901 DT Patent LA German
- AB Propane-1,3-diol is manufd. from a glycerol-contg. soln. (5-20% by wt.) with a microorganism such as *Clostridium*, *Enterobacterium*, *Lactobacillus*, "Bacillus", *Citrobacter*, or *Klebsiella* in a yield of, gloreq.0.5 g/mL. *Klebsiella pneumoniae* DSM 2026 was batch-cultured at 37 degree. under anaerobic conditions to yield a max. of 2.3 g propane-1,3-diol from a starting glycerol concn. Of 100 g/L; other glycerol concns. (50-200 g/L) produced lower yields.
- L13 ANSWER 8 OF 8 CAPLUS COPYRIGHT 1998 ACS
- AN 1993:214106 CAPLUS DN 98:214106
- TI Neutral solvent production from halophilic, photolithotrophically grown algae by linked fermentations
- AU Nakas, J. P.; Schaedle, M.; Parkinson, C. M.; Coonley, C. E.; Tanenbaum, S. W.
- CS Coll. Environ. Sci. For. SUNY, Syracuse, NY 13210 USA
- SO Comm. Eur. Communities, [Rep.] EUR (1993), EUR 8245, Energy Biomass, 298-302 CODEN: CECEED9 DT Report LA English
- AB Five species of *Dunaliella* were examd. for glycerol [56-81-5] accumulation, growth rate, cell d., and protein and chlorophyll content. The suitability of each algal species for such bioconversions was judged according to glycerol accumulation and quantities of neutral solvents produced after sequential bacterial fermentns. When grown in 2M NaCl, with 24 mM NaHCO<sub>3</sub> or 3%
- CO<sub>2</sub> at 28 degree., and with 25,000 lx at container surface, 4 of the 5 species tested (*D. tertiolecta*, *D. parva*, and *D. bardawilii*) produced 10-20 mg of glycerol/L. A *Clostridium* converted an algal biomass mixt. supplemented with 4% glycerol to approx. 18 g/L of mixed alcs. [EtOH (64-17-5), 1,3-propanediol (504-63-2), and BuOH (71-36-3)]. Acetone was not detected. A soil isolate, tentatively classified as a member of the genus "Bacillus", converts glycerol into EtOH at a final concn. of 7.0-9.6 g/L. An enrichment culture from sewage sludge resolved to contain 2 gram-neg. rods converts the algal biomass-glycerol mixt. solely to 1,3-propanediol (504-63-2) at a final concn. of 4.2-5.3 g/L. Addnl., *Dunaliella* concs., of ltoreq.200-fold, can be directly fermented to mixed solvents.